

Positive Quality Intervention: Acalabrutinib (Calquence®) In Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma

Description:

- The purpose of this PQI is to discuss clinical considerations around the use of acalabrutinib to optimize outcomes for patients with chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL).

Background:

- Acalabrutinib is a selective, covalent Bruton tyrosine kinase (BTK) inhibitor that irreversibly binds BTK, blocking B-cell receptor signaling. This inhibits downstream pathways that promote CLL cell proliferation, survival, adhesion, and migration.
- FDA approved indications include:
 - CLL/SLL
 - As monotherapy
 - In combination with obinutuzumab or venetoclax in patients with previously untreated disease
 - Mantel cell lymphoma (MCL)
 - As monotherapy for the treatment of adult patients with MCL who have received at least one prior therapy
 - In combination with bendamustine and rituximab for the treatment of adult patients with previously untreated MCL who are ineligible for autologous hematopoietic stem cell transplantation
- The most common adverse reactions ($\geq 30\%$), excluding laboratory abnormalities, are upper respiratory tract infection, diarrhea, headache, and musculoskeletal pain.
- The most common Grade 3 or 4 laboratory abnormalities ($\geq 10\%$) are absolute neutrophil count decreased, uric acid increased, absolute lymphocyte count decreased, and platelets decreased.

PQI Process:

Upon the receipt of a new prescription of acalabrutinib for CLL/SLL:

- Verify dose: acalabrutinib 100 mg by mouth every 12 hours, taken whole with water, with or without food
 - If dose is missed by > 3 hours, skip and take the next at its regularly scheduled time
 - Avoid in patients with severe hepatic impairment
 - No dose adjustment needed in mild to moderate hepatic or renal impairment (use in severe renal impairment or with dialysis has not yet been evaluated)
 - In combination with obinutuzumab
 - Start acalabrutinib at Cycle 1 (each cycle is 28 days)
 - Start obinutuzumab at Cycle 2 for a total of 6 cycles (refer to obinutuzumab prescribing information for recommended dosing)
 - Administer acalabrutinib **prior** to obinutuzumab when given on the same day
 - Administers until disease progression or unacceptable toxicity
 - In combination with venetoclax

IMPORTANT NOTICE: NCODA has developed this Positive Quality Intervention platform. This platform is intended as an educational aid, does not provide individual medical advice, and does not substitute for the advice of a qualified healthcare professional. This platform does not cover all existing information related to the possible uses, directions, doses, precautions, warning, interactions, adverse effects, or risks associated with the medication. The materials contained in this platform do not constitute or imply endorsement, recommendation, or favoring of this medication by NCODA. NCODA does not ensure the accuracy of the information presented and assumes no liability relating to its accuracy. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional. It is the individual's sole responsibility to seek guidance from a qualified healthcare professional. *Updated 10.2026 PQI-165*

- Start acalabrutinib at Cycle 1 (each cycle is 28 days)
 - Start venetoclax at Cycle 3 for total of 12 cycles; starting dose venetoclax is 20 mg daily for first week of treatment and increases weekly as per dosing schedule for 5-week ramp up (up to 400 mg daily) as described in the venetoclax PI.
 - Continue treatment with acalabrutinib + venetoclax until disease progression, unacceptable toxicity or completion of 14 cycles of treatment
- Review patient medication list for possible drug-drug interactions
 - Strong CYP3A4 inducer: avoid use; if use cannot be avoided, increase dosage to 200 mg every 12 hours
 - Strong CYP3A4 inhibitor: avoid use; if the inhibitor is a short-term medication (e.g., anti-infectives for up to seven days), stop acalabrutinib and resume after at least 24 hours post inhibitor discontinuation
 - Moderate CYP3A4 inhibitor: reduce acalabrutinib dosage to 100 mg daily
 - Tablet formulation may be co-administered with gastric reducing agents
- Adverse Event Monitoring and Management
 - Infections, including opportunistic infections
 - Monitor for signs/symptoms of infections and treat promptly
 - Consider prophylaxis in patients who are at increased risk for opportunistic infections
 - Hemorrhagic events
 - Monitor for bleeding and manage appropriately
 - Consider the benefit-risk of withholding acalabrutinib for 3-7 days pre- and post-surgery depending on type of surgery and the risk of bleeding
 - Caution in patients on antithrombotic agents
 - Myelosuppression (neutropenia, anemia, thrombocytopenia, and lymphopenia)
 - Monitor complete blood cell counts regularly
 - Cardiac arrhythmia
 - Monitor for symptoms of arrhythmias (ex. palpitations, dizziness, syncope, dyspnea) and manage appropriately
 - Risk of arrhythmias may be increased in patients with cardiac risk factors, hypertension, previous arrhythmias, and acute infection
 - Skin cancer as secondary malignancy
 - Monitor patients for skin cancer and advise protection from sun exposure
 - Hepatotoxicity, including drug-induced liver injury (DILI)
 - Evaluate bilirubin and transaminases at baseline and throughout treatment
 - For patients who develop abnormal liver tests, monitor more frequently for liver test abnormalities and clinical signs and symptoms of hepatic toxicity
 - If DILI is suspected, withhold acalabrutinib; upon confirmation of DILI, discontinue acalabrutinib

Table 1. Recommended Dosage Modifications for Adverse Reactions (acalabrutinib monotherapy/in combination with obinutuzumab*)

Event	Adverse Reaction Occurrence	Dosage Modification (Starting dose = 100 mg approximately every 12 hours)
Grade 3 or greater nonhematologic toxicities, Grade 3 thrombocytopenia with bleeding, Grade 4 thrombocytopenia or Grade 4 neutropenia lasting longer than 7 days	First and Second	Interrupt acalabrutinib. Once toxicity has resolved to Grade 1 or baseline level, acalabrutinib may be resumed at normal starting dose.
	Third	Interrupt acalabrutinib. Once toxicity has resolved to Grade 1 or baseline level, acalabrutinib may be resumed at a reduced frequency of 100 mg once daily.
	Fourth	Discontinue acalabrutinib.

*See [CALQUENCE PI Table 4](#) for recommended dosage modifications when given in combination with venetoclax

Patient-Centered Activities:

- Patient Education
 - Provide Patient Education Sheet (PES) and review with patient
 - [Acalabrutinib PES](#)
 - [Acalabrutinib + Venetoclax PES](#)
 - Instruct patients to report any signs or symptoms of atrial fibrillation or flutter such as palpitations, dizziness, faint feeling, or chest discomfort
 - Counsel patients to report any signs or symptoms of infections including new or worsening cough or shortness of breath, sore throat, chills, notably increased fatigue, or fever of 100.4° F or higher
 - Counsel patients to report any new or uncontrolled bleeding including nosebleed lasting > 5 minutes despite pressure, a cut that won't stop bleeding, heavy gum bleeding when brushing or flossing, or blood in urine or stool
 - Patients should be made aware of the increased bleeding risk associated with acalabrutinib
 - Due to this risk, they may need to hold their medication prior to any procedures
 - Ensure patients have access to supportive care medications for diarrhea such as loperamide
- Patient Assistance: [CALQUENCE Co-Pay Savings Program](#) & [AstraZeneca Access 360](#)

References:

1. [CALQUENCE® \(acalabrutinib\) \[prescribing information\]. Wilmington, DE: AstraZeneca Pharmaceuticals LP; 2026.](#)
2. Sharman JP, et al. Acalabrutinib with or without obinutuzumab versus chlorambucil and obinutuzumab for treatment-naïve chronic lymphocytic leukemia (ELEVATE TN): a randomized, controlled, phase 3 trial. *Lancet*. 2020 Apr 18;395(10232):1278-1291.
3. Ghia P, et al. ASCEND: Phase III, Randomized Trial of Acalabrutinib Versus Idelalisib Plus Rituximab or Bendamustine Plus Rituximab in Relapsed or Refractory Chronic Lymphocytic Leukemia. *J Clin Oncol*. 2020 Sept 01;38(25):2849-2861.
4. Byrd J, et al. Acalabrutinib Versus Ibrutinib in Previously Treated Chronic Lymphocytic Leukemia: Results of the First Randomized Phase III Trial. *J Clin Oncol*. 2021 Nov 01;39(31):3441-3452.