

Brentuximab Vedotin

Care Team Contact Information: _____

Pharmacy Contact Information: _____

Diagnosis: _____

- This treatment is often used for lymphoma, but it may also be used for other diagnoses.

Goal of Treatment: _____

- Treatment may continue for a certain time period, until it no longer works, or until side effects are no longer controlled.

Treatment Regimen

Treatment Name	How the Treatment Works	How the Treatment is Given
Brentuximab Vedotin (bren-TUK-sih-mab veh-DOH-tin): Adcetris (ad-SEH-tris)	Stops cancer cell growth by targeting a protein and delivering medicine directly to the cancer cell.	Infusion given through a vein.

Treatment Administration and Schedule: Treatment is typically repeated every 3 weeks. This length of time is called a “cycle”.

- Brentuximab vedotin may be given with other treatments. Talk with your care team about your exact treatment.

Treatment Name	Cycle 1							Next Cycle
	Day 1	Day 2	Day 3	Day 4	Day 5	...	Day 21	Day 1
Brentuximab Vedotin	✓							✓

Appointments: Appointments may include regular check-ups with your care team, treatment appointments, lab visits, and imaging tests. It's important to keep your appointments whenever you can. If you miss any appointments, call your care provider as soon as possible to reschedule your appointment.

Supportive Care to Prevent and Treat Side Effects

Description	Supportive Care Given at the Clinic or Hospital	Supportive Care Taken at Home
To help prevent or treat nausea or vomiting	<hr/> <hr/> <hr/>	<hr/> <hr/> <hr/>
To help your body make white blood cells to fight infections	<hr/> <hr/> <hr/>	<hr/> <hr/> <hr/>
To help prevent infusion-related reactions	<hr/> <hr/> <hr/>	<hr/> <hr/> <hr/>

Common Side Effects

Side Effect	Important Information
<p>Low White Blood Cell (WBC) Count and Increased Risk of Infection</p>	<p>Description: WBCs help protect the body against infections. If you have a low WBC count, you may be at a higher risk of infection.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Wash your hands and bathe regularly. • Avoid crowded places. • Stay away from people who are sick. • Your care team may prescribe a drug that promotes the growth of WBCs. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • Fever of 100.4 °F (38°C) or higher • Chills • Cough • Sore throat • Painful urination • Tiredness that is worse than normal • Skin infections (red, swollen, or painful areas)
<p>Low Platelet Count</p>	<p>Description: Platelets help the blood clot and heal wounds. If you have low platelet counts, you are at a higher risk of bruising and bleeding.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Blow your nose gently and avoid picking it. • Brush your teeth gently with a soft toothbrush and maintain good oral hygiene. • Use an electric razor for shaving and a nail file instead of nail clippers. • Avoid over-the-counter medications that may increase the risk of bleeding, such as NSAIDs. • Talk with your care team or dentist before medical or dental procedures, as you may need to pause your treatment. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • Nosebleed lasting over 5 minutes despite pressure • Cut that continues to bleed • Significant gum bleeding when flossing or brushing • Severe headaches • Blood in your urine or stool • Blood in your spit after a cough
<p>Low Red Blood Cell (RBC) Count and Hemoglobin (Hgb)</p>	<p>Description: RBCs and Hgb help bring oxygen to your body's tissues and take away carbon dioxide. If you have low RBC counts or Hgb, you may feel weak, tired, or look pale.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Get 7 to 8 hours of sleep each night. • Avoid operating heavy machinery when tired. • Balance work and rest, staying active but resting when needed. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • Shortness of breath • Dizziness • Fast or abnormal heartbeats • Severe headache

<p>Fatigue</p>	<p>Description: Fatigue is a constant and sometimes strong feeling of tiredness.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Routine exercise has been shown to decrease levels of fatigue. Work with your care team to find the right type of exercise for you. • Ask your family and friends for help with daily tasks and emotional support. • Try healthy ways to feel better, like meditation, writing in a journal, doing yoga, and using guided imagery to lower anxiety and feel good. • Make a regular sleep schedule and limit naps during the day so you can sleep better at night, aiming for 7 to 8 hours of sleep. • Don't use heavy machines or do things that need your full attention if you're very tired to avoid accidents. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • Tiredness that affects your daily life • Tiredness all the time, and it doesn't get better with rest • Dizziness and weakness, along with being tired
<p>Nausea and Vomiting</p>	<p>Description: Nausea is an uncomfortable feeling in your stomach or the need to throw up. This may or may not cause vomiting.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Eat smaller, more frequent meals. • Avoid fatty, fried, spicy, or highly sweet foods. • Eat bland foods at room temperature and drink clear liquids. • If you vomit, start with small amounts of water, broth, or other clear liquids when you are ready to eat again. If that stays down, then try soft foods (such as gelatin, plain cornstarch pudding, yogurt, strained soup, or strained cooked cereal). Slowly work up to eating solid food. • Your care team may prescribe medicine for these symptoms. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • Vomiting for more than 24 hours • Vomiting that's nonstop • Signs of dehydration (like feeling very thirsty, having a dry mouth, feeling dizzy, or having dark urine) • Blood or coffee-ground-like appearance in your vomit • Bad stomach pain that doesn't go away after vomiting

<p>Diarrhea</p>	<p>Description: Diarrhea is when you have loose, watery bowel movements more often than usual. The need to use the bathroom may occur urgently.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Keep track of how many times you go to the bathroom each day. • Drink 8 to 10 glasses of water or other fluids every day, unless your doctor tells you otherwise. • Eat small meals of mild, low-fiber foods like bananas, applesauce, potatoes, chicken, rice, and toast. • Avoid eating foods with high fiber (like raw vegetables, fruits, and whole grains), foods that cause gas (like broccoli and beans), dairy foods (like yogurt and milk), and spicy, fried, and greasy foods. • Your care team may recommend medicine (such as loperamide) for diarrhea. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • 4 or more bowel movements than normal in 24 hours • Dizziness or lightheadedness while having diarrhea • Bloody diarrhea
<p>Numbness or Tingling in the Hands and Feet</p>	<p>Description: Nerve pain and tingling are uncomfortable sensations caused by nerve damage or irritation. Pain may be sharp, burning, or deep, while tingling feels like pins-and-needles or mild electric shocks, often in the hands, feet, arms, or legs.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Track your pain levels, sensations, and any triggers or factors that make the pain worse or better. • Check your feet daily for any injuries or changes, especially if you have numbness or tingling that affects your feeling. • Your care team may prescribe or recommend medicine for symptoms. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • Pins and needles” or burning feeling in your hands or feet • Trouble moving your arms or legs • Trouble keeping your balance

<p>Muscle or Joint Pain</p>	<p>Description: Muscle pain feels like soreness, aches, cramps, or stiffness in one or more muscles. It may also include tenderness or weakness. Joint pain happens where two bones come together and can feel sharp, dull, throbbing, or burning. It often causes stiffness, swelling, and difficulty moving.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> Track your pain levels, areas of discomfort, and any activities that worsen or improve your symptoms. Engage in gentle exercises like walking, stretching, or yoga to maintain mobility and strength, but consult your care team before starting any new exercise routine. Apply a warm compress to relax stiff muscles or use cold packs to reduce swelling and numb pain in affected areas. Your care team may prescribe or recommend medications, including over-the-counter pain relievers. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> Uncontrolled muscle or joint pain Swelling, redness, or warmth in the joints Weakness Difficulty walking or moving
<p>Rash or Itchy Skin</p>	<p>Description: Rash or itching skin can cause symptoms like redness, swelling, and varying bumps, such as small red spots, welts, or patches.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> Take a warm (not hot) bath or shower every day with mild, unscented soap or body wash. Pat skin dry (instead of rubbing) with a towel. Use unscented lotion or moisturizing cream immediately after shower to prevent cracking. Use lip balm for dry or chapped lips. Avoid activities that put you at risk for cuts, injuries or burns. Use an electric shaver, not a razor, to prevent cuts. Rinse cuts and scrapes immediately under running water and remove any debris from the wound. Wash the skin around the wound with soap. To avoid irritation, do not use soap on the wound. If you have a low platelet count and are at risk for bleeding issues, ask your doctor how to care for cuts. Avoid direct sun when possible. Wear a broad-spectrum sunscreen with SPF 30 or higher and reapply every 2 hours. Use unscented, gentle laundry detergent. Some laundry detergents can irritate sensitive skin. Your care team may recommend medicine for skin problems. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> Skin rash Raised red bumps Redness of the skin Very dry skin that may affect the mucous membranes (such as mouth and eyes)

Select Rare Side Effects

Side Effect	Talk to Your Care Team if You Have Any of These Signs or Symptoms	
Progressive Multifocal Leukoencephalopathy (PML) (Boxed Warning)	<p>PML is a rare, serious brain infection caused by a virus that can happen in people who receive this treatment. People with weakened immune systems can get PML. PML can result in death or severe disability. There is no known treatment, prevention, or cure for PML.</p> <ul style="list-style-type: none"> • Confusion • Dizziness or loss of balance • Difficulty walking or talking • Decreased strength or weakness on one side of your body • Vision problems 	
Tumor Lysis Syndrome	<p>Tumor lysis happens when cancer cells break apart and flood your bloodstream with chemicals and toxins faster than your body can get rid of them. TLS is a group of conditions that affect your heart, kidneys, and muscles.</p> <ul style="list-style-type: none"> • Fever of 100.4°F (38°C) or higher • Chills • Nausea • Vomiting • Confusion • Shortness of breath • Seizures • Irregular heartbeat • Dark or cloudy urine • Unusual tiredness • Muscle or joint pain 	
Stomach and Serious Bowel Problems	<ul style="list-style-type: none"> • Severe stomach-area (abdomen) pain • Repeated vomiting during treatment 	
Liver Problems	<ul style="list-style-type: none"> • Yellowing of your skin or the whites of your eyes • Severe nausea or vomiting, pain on the right side of your stomach area (abdomen) • Dark urine (tea colored) • Bleeding or bruising more easily than normal 	
Lung Problems	<ul style="list-style-type: none"> • Cough • Shortness of breath • Chest pain 	
High Blood Sugar	<ul style="list-style-type: none"> • Frequent urination • Drowsiness • Increased thirst • Loss of appetite • Blurred vision • Fruity smell on your breath • Confusion • Nausea, vomiting, or stomach pain • It becomes harder to control your blood sugar 	
Severe Skin Reactions	<ul style="list-style-type: none"> • A red, itchy, skin rash • Peeling of your skin or blisters • Severe itching • Fever of 100.4°F (38°C) or higher 	
Infusion Reaction	<ul style="list-style-type: none"> • Chills or shaking • Itching, rash, or flushing • Trouble breathing or wheezing; tongue-swelling • Dizziness or feeling faint • Fever of 100.4°F (38°C) (or 38°C) or higher • Pain in your back or neck 	

**Before starting treatment, ask your care team when to call 9-1-1 or seek emergency help.
If you experience any new, worsening, or uncontrolled side effects, contact your care team immediately.**

Intimacy, Fertility, Pregnancy, and Breastfeeding

- Treatment may **change how you feel about intimacy and your body**. However, physical closeness—such as holding hands and hugging—remains safe. It is common to have questions about intimacy. If needed, talk to your care team for guidance.
- Treatment may affect your **ability to have children**. It may damage your reproductive organs or stop them from working. If you are worried about fertility, talk to your care team before starting treatment.
- Treatment may **harm an unborn baby**.
 - If you are able to become pregnant, take a pregnancy test before starting treatment.
 - Use an effective method of birth control during treatment and for 2 months after your last dose.
 - If you think you might be pregnant or if you become pregnant, tell your care team right away.
 - If your partner(s) could become pregnant, use an effective method of birth control—such as condoms—during treatment and for 4 months after your last dose.
- **Do NOT breastfeed** during treatment.

Handling Body Fluids and Waste

Some drugs you receive may stay in your urine, stool, sweat, or vomit for many days after treatment. Because many cancer drugs are toxic, your body waste may also be dangerous to touch. To help protect yourself, your loved ones, and the environment, **follow these instructions** for at least **3 weeks** after each dose of **brentuximab vedotin**:

- People who are pregnant should avoid touching anything that may be soiled with body fluids from the patient.
- You can use your usual toilet. Always close the lid and flush to discard all waste. If you have a low-flow toilet, flush twice.
- If the toilet or seat is soiled with urine, stool, or vomit, clean the surface after each use before others use it.
- Wash your hands with soap and water for at least 20 seconds after using the toilet.
- If you need a bedpan, inform your caregiver so they can wear gloves and assist with cleanup. Wash the bedpan with soap and water daily.
- If you cannot control your bladder or bowels, use a disposable pad with a plastic back, a diaper, or a sheet to absorb waste.
- Wash any skin exposed to body waste with soap and water.
- Wash soiled linens or clothing separately from other laundry. If you don't have a washer, place them in a plastic bag until they can be washed.
- Wash your hands with soap and water after touching soiled linens or clothing.

Additional Information

- **Tell your care team about all the medicines you take.**
This includes prescriptions, over-the-counter drugs, vitamins, and herbal products. Before starting any new medicine, supplement, or vaccine, ask your care team first.
- **This Patient Education Sheet may not describe all possible side effects.**
Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Notes

Updated Date: December 17, 2025

Scan the QR code below to access this education sheet.



Important notice: The Association of Cancer Care Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), Network for Collaborative Oncology Development & Advancement, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated in gathering information for and developing this patient education guide. This guide represents a brief summary of the medication derived from information provided by the drug manufacturer and other resources.

This guide does not cover all existing information related to the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this medication and should not substitute for the advice of a qualified healthcare professional. Provision of this guide is for informational purposes only and does not constitute or imply endorsement, recommendation, or favoring of this medication by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional.

Permission: Patient Education Sheets are provided as a free educational resource for patients with cancer and their caregivers in need of concise, easy to understand information about cancer therapy. Healthcare providers are permitted to copy and distribute the sheets to patients as well as direct patients to the Patient Education Sheets website. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holders. Permission requests, including direct linking from Electronic Health Records, and licensing inquiries should be emailed to patienteducation sheets@ncoda.org.

Copyright © 2025 by Network for Collaborative Oncology Development & Advancement, Inc. All rights reserved.

PES-383