Lomustine

Patient Education Sheets
Understanding Your Cancer Treatment

Care Team Contact Information: _____

Pharmacy Contact Information: _____

Diagnosis: _____

- This treatment is used for brain tumors (cancer) and Hodgkin lymphoma (HL).
- It may also be used for other reasons.

Goal of Treatment: _____

- Treatment may continue for a certain time period, until it no longer works, or until side effects are no longer controlled.

Treatment Regimen

Treatment Name	How the Treatment Works	How the Treatment is Given
Lomustine (loh-MUS-teen): Gleostine (GLEE-oh-steen)	Slows down or stops the growth of cancer cells by damaging the genetic material that cancer cells need to multiply.	Capsule(s) taken by mouth.

Treatment Administration and Schedule

Treatment is typically repeated every 6 weeks. This length of time is called a "cycle."

Your lomustine dosing instructions:

_____

_____

_____

- Lomustine comes in 3 capsule strengths: 10 mg, 40 mg, and 100 mg. Your care team will tell you which capsules to take.
- Your dose is based on many factors, including your height and weight, overall health, and diagnosis.
- Take lomustine once every 6 weeks.
- Swallow lomustine capsules whole. Do not break, crush, or chew lomustine tablets.
- Take lomustine with or without food. Taking lomustine without food may lower the risk of nausea and vomiting.
- Do not change your dose or stop taking lomustine unless your care team tells you to.
- Your care team may tell you to change your dose, temporarily stop, or completely stop taking lomustine if you develop certain side effects.
- If you miss a dose of lomustine, or if you vomit after taking your lomustine, take your next dose at the next scheduled time. Do not take another dose on the same day.
- If you take too much lomustine, call your care team or go to the nearest hospital emergency room right away.

Brought to you by: ACCC Association of Cancer Care Centers, HOPA Hematology/Oncology Pharmacy Association, NCODA Network for Collaborative Oncology Development & Advancement, ONS Oncology Nursing Society

Treatment Administration and Schedule (Continued)

- Lomustine is taken on Day 1.

Treatment Name	Cycle 1								Next Cycle
	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	…	Day 42	Day 1
Lomustine	✓								✓

Storage and Handling of Lomustine

- Store lomustine at room temperature between 68°F and 77°F (20ºC and 25ºC).
- If someone else is giving you your medication, they should wear gloves or pour the pills directly from the bottle into the cap, a small cup, or your hand without touching them. They should wash their hands before and after handling the pills.
- Keep lomustine and all medicines out of the reach of children and pets.
- Ask your care team how to safely throw away any unused lomustine.

Appointments

Appointments may include regular check-ups with your care team, lab visits, and imaging tests. It's important to keep your appointments whenever you can. If you miss an appointment, call your care team as soon as possible to reschedule it.

Supportive Care to Prevent and Treat Side Effects

Description	Supportive Care Taken at Home
To help prevent or treat nausea and vomiting	
Other	

Brought to you by:

ACCC Association of Cancer Care Centers™

HOPA Hematology/Oncology Pharmacy Association

NCODA Network for Collaborative Oncology Development & Advancement

ONS Oncology Nursing Society

Common Side Effects

Side Effect	Important Information
Low White Blood Cell (WBC) Count (Neutropenia) and Increased Risk of Infection (Boxed Warning)	Description: WBCs help protect your body from infections. A low WBC count increases your risk of infection. Recommendations: • Wash your hands often and bathe regularly. • Avoid crowded places and close contact with people who are sick. • Follow food safety and wound care advice from your care team. Talk to your care team if you have: • Fever of 100.4°F (38°C) or higher • Chills • New or worsening cough or sore throat • Painful urination or signs of a urinary infection • Feeling much more tired than usual • Red, swollen, warm, or painful areas on the skin
Low Platelet Count (Thrombocytopenia) (Boxed Warning)	Description: Platelets help your blood clot and wounds heal. A low platelet count increases your risk of bruising and bleeding. Recommendations: • Blow your nose gently and avoid picking it. • Brush your teeth gently with a soft toothbrush and keep good oral hygiene. • Use an electric razor for shaving and a nail file instead of nail clippers. • Avoid over-the-counter medicines that can increase bleeding risk (for example, nonsteroidal anti-inflammatory drugs (NSAIDs) like ibuprofen). • Talk with your care team or dentist before medical or dental procedures — you may need to pause treatment. Talk to your care team if you have: • A nosebleed lasting more than 5 minutes despite pressure • A cut that continues to bleed • Heavy gum bleeding when brushing or flossing • Sudden or severe headache • Blood in your urine or stool • Blood in your spit after coughing
Low Red Blood Cell (RBC) Count and Hemoglobin (Hgb) (Anemia) (Boxed Warning)	Description: Red blood cells carry oxygen throughout your body. Low levels can make you feel tired, weak, dizzy, or short of breath. Recommendations: • Aim for 7 to 8 hours of sleep each night. • Do not drive, operate heavy machinery, or do other dangerous activities if you are very tired. • Balance activity and rest — stay as active as you can, but rest when needed. • Eat a balanced diet and follow any nutrition or supplement advice from your care team. Talk to your care team if you have: • Shortness of breath • Dizziness or fainting • Fast or irregular heartbeats • Sudden or severe headache

Brought to you by: ACCC Association of Cancer Care Centers, HOPA Hematology/Oncology Pharmacy Association, NCODA Network for Collaborative Oncology Development & Advancement, ONS Oncology Nursing Society

Nausea and Vomiting	Description: Nausea is an uncomfortable feeling in your stomach or the need to throw up. You may or may not vomit. Recommendations: • Eat smaller, more frequent meals. • Avoid fatty, fried, spicy, or highly sweet foods. • Eat bland foods at room temperature and drink clear liquids. • If you vomit, start with small sips of water, broth, or other clear liquids. If these stay down, try soft foods (such as gelatin, plain cornstarch pudding, yogurt, strained soup, or strained cooked cereal) and gradually return to solid foods. • Your care team may prescribe medicine for these symptoms. Talk to your care team if you have: • Vomiting for more than 24 hours • Nonstop vomiting • Signs of dehydration (very thirsty, dry mouth, dizziness, or dark urine) • Blood in your vomit or vomit that looks like coffee grounds • Severe stomach pain that does not go away after vomiting

Brought to you by:
ACCC Association of Cancer Care Centers
HOPA Hematology/Oncology Pharmacy Association
NCODA Network for Collaborative Oncology Development & Advancement
ONS Oncology Nursing Society

Select Rare Side Effects

Side Effect	Talk to Your Care Team if You Have Any of These Signs or Symptoms	
Liver Problems	• Yellowing of your skin or the white part of your eyes (jaundice) • Severe nausea or vomiting	• Pain on the right side of your stomach area (abdomen) • Dark, tea-colored urine • Bleeding or bruising more easily than normal
Kidney Problems	• Decrease in your amount of urine • Blood in your urine	• Swelling of your ankles • Loss of appetite
Lung Problems	• Cough • Shortness of breath	• Chest pain or tightness
Risk of New Cancers	There is a risk of developing one or more new cancers during or after treatment. Talk with your care team about this risk, and ask about the signs and symptoms to watch for.	

Before starting treatment, ask your care team when to call 9-1-1 or seek emergency help.
If you experience any new, worsening, or uncontrolled side effects, contact your care team immediately.

Intimacy, Fertility, Pregnancy, and Breastfeeding

- Treatment may **change how you feel about intimacy and your body.** However, physical closeness—such as holding hands and hugging—remains safe. It is common to have questions about intimacy. If needed, talk to your care team for guidance.
- Treatment can affect your **ability to have children**. It may damage your reproductive organs or stop them from working. If you are worried about fertility, talk to your care team before starting treatment.
- Treatment may **harm an unborn baby**.
    - If you are able to become pregnant:
        - Your care team will verify your pregnancy status before treatment.
        - Use an effective method of birth control during treatment and for 2 weeks after your last dose of lomustine.
        - If you think you might be pregnant or if you become pregnant, tell your care team right away.
    - If your partner is able to become pregnant, use an effective method of birth control—such as condoms—during treatment and for 4 months after your last dose of lomustine.
- **Do not breastfeed** during treatment and for 2 weeks after your last dose of lomustine.

Brought to you by: ACCC Association of Cancer Care Centers, HOPA Hematology/Oncology Pharmacy Association, NCODA Network for Collaborative Oncology Development & Advancement, ONS Oncology Nursing Society

Handling Body Fluids and Waste

Some drugs you receive may stay in your urine, stool, sweat, or vomit for many days after treatment. Because many cancer drugs are toxic, your body waste may also be dangerous to touch. To help protect yourself, your loved ones, and the environment, **follow these instructions** for at least **6 days** after each dose of **lomustine**:

- People who are pregnant should avoid touching anything that may be soiled with body fluids from the patient.
- You can use your usual toilet. Always close the lid and flush to discard all waste. If you have a low-flow toilet, flush twice.
- If the toilet or seat is soiled with urine, stool, or vomit, clean the surface after each use before others use it.
- Wash your hands with soap and water for at least 20 seconds after using the toilet.
- If you need a bedpan, inform your caregiver so they can wear gloves and assist with cleanup. Wash the bedpan with soap and water daily.
- If you cannot control your bladder or bowels, use a disposable pad with a plastic back, a diaper, or a sheet to absorb waste.
- Wash any skin exposed to body waste with soap and water.
- Wash soiled linens or clothing separately from other laundry. If you don't have a washer, place them in a plastic bag until they can be washed.
- Wash your hands with soap and water after touching soiled linens or clothing.

Additional Information

- **Tell your care team about all the medicines you take**.
  This includes prescriptions, over-the-counter drugs, vitamins, and herbal products. Before starting any new medicine, supplement, or vaccine, ask your care team first.
- **This Patient Education Sheet may not describe all possible side effects.**
  Call your care team for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Notes

Updated Date: November 5, 2026

Brought to you by:
ACCC ASSOCIATION OF CANCER CARE CENTERS™
HOPA Hematology/Oncology Pharmacy Association
NCODA Network for Collaborative Oncology DEVELOPMENT & ADVANCEMENT
ONS Oncology Nursing Society

Scan the QR code below to access this education sheet.

Important Notice:
The Association of Cancer Care Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), Network for Collaborative Oncology Development & Advancement, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated to gather information and develop this patient education guide. This guide is a summary of the medication, based on information provided by the drug manufacturer and other sources.

This guide does not cover all information available on the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this medication. It should not be used as a substitute for the advice of a qualified healthcare professional. The provision of this guide is for informational purposes only. It does not constitute or imply endorsement, recommendation, or favoring of this medication by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. All decisions regarding this medication should be made with the guidance and direction of a qualified healthcare professional.

Permission:
Patient Education Sheets are provided as a free educational resource for patients with cancer and their caregivers in need of concise, easy-to-understand information about cancer therapy. Healthcare providers may copy and distribute the sheets to patients and direct them to the Patient Education Sheets website. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holders. Permission requests, including direct linking from Electronic Health Records, and licensing inquiries should be emailed to patienteducationsheets@ncoda.org.

Copyright © 2026 by Network for Collaborative Oncology Development & Advancement, Inc. All rights reserved.

PES-260

Brought to you by:
ACCC Association of Cancer Care Centers
HOPA Hematology/Oncology Pharmacy Association
NCODA Network for Collaborative Oncology Development & Advancement
ONS Oncology Nursing Society