Selinexor

Care Team Contact Information: _____

Pharmacy Contact Information: _____

Diagnosis: _____

- This treatment is used for multiple myeloma (MM).
- It may also be used for other reasons.

Goal of Treatment: _____

- Treatment may continue for a certain time period, until it no longer works, or until side effects are no longer controlled.

Treatment Regimen

Treatment Name	How the Treatment Works	How the Treatment is Given
Selinexor (SEH-lih-NEK-sor) Xpovio (ex-POH-vee-oh)	Slows down or stops the growth of cancer cells by blocking a specific protein that helps them survive.	Tablet(s) taken by mouth.

Treatment Administration and Schedule

- Treatment is typically repeated once or twice every week.
- Selinexor is often given in combination with other treatments. Talk with your care team about your exact treatment and schedule.

Your selinexor dosing instructions:

_____

_____

_____

- Selinexor comes in 6 tablet strengths: 10 mg, 20 mg, 40 mg, 50 mg, 60 mg, and 80 mg.
- Your dose is based on many factors, including your overall health and diagnosis.
- Swallow selinexor tablets whole with water. Do not break, chew, crush, or divide the tablets.
- It is important for you to drink enough fluids to help prevent dehydration and to eat enough calories to help prevent weight loss during treatment with selinexor. Talk to your care team if this is a problem for you.
- Do not change your dose or stop taking selinexor unless your care team tells you to.
- Your care team may tell you to change your dose, temporarily stop, or completely stop taking selinexor if you develop certain side effects.
- If you miss a dose of selinexor, take your next dose at your next regularly scheduled day and time.

Brought to you by: ACCC Association of Cancer Care Centers, HOPA Hematology/Oncology Pharmacy Association, NCODA Network for Collaborative Oncology Development & Advancement, ONS Oncology Nursing Society

Treatment Administration and Schedule (Continued)

- If you vomit after taking a dose of selinexor, do not take an extra dose. Take your next dose at your next regularly scheduled day and time.
- If you take too much selinexor, call your care team or go to the nearest hospital emergency room right away.

Storage and Handling of Selinexor

- Store selinexor at or below 86°F (30°C).
- Selinexor comes in a child-resistant blister pack.
- Keep selinexor and all medicines out of the reach of children and pets.
- Ask your care team how to safely throw away any unused selinexor.

Appointments

Appointments may include regular check-ups with your care team, lab visits, and imaging tests. It's important to keep your appointments whenever you can. If you miss an appointment, call your care team as soon as possible to reschedule it.

Supportive Care to Prevent and Treat Side Effects

Description	Supportive Care Taken at Home
To help prevent or treat nausea and vomiting	
Other	

Brought to you by:

ACCC Association of Cancer Care Centers™

HOPA Hematology/Oncology Pharmacy Association

NCODA Network for Collaborative Oncology Development & Advancement

ONS Oncology Nursing Society

Common Side Effects

Side Effect	Important Information
Low White Blood Cell (WBC) Count (Neutropenia) and Increased Risk of Infection	Description: WBCs help protect your body from infections. A low WBC count increases your risk of infection. Recommendations: • Wash your hands often and bathe regularly. • Avoid crowded places and close contact with people who are sick. • Follow food safety and wound care advice from your care team. • Your care team may prescribe medicine to help your WBCs recover. Talk to your care team if you have: • Fever of 100.4°F (38°C) or higher • Chills • New or worsening cough or sore throat • Painful urination or signs of a urinary infection • Feeling much more tired than usual • Red, swollen, warm, or painful areas on the skin (possible skin infection)
Low Platelet Count (Thrombocytopenia)	Description: Platelets help your blood clot and wounds heal. A low platelet count increases your risk of bruising and bleeding. Recommendations: • Blow your nose gently and avoid picking it. • Brush your teeth gently with a soft toothbrush and keep good oral hygiene. • Use an electric razor for shaving and a nail file instead of nail clippers. • Avoid over-the-counter medicines that can increase bleeding risk (for example, nonsteroidal anti-inflammatory drugs (NSAIDs) like ibuprofen). • Talk with your care team or dentist before medical or dental procedures — you may need to pause treatment. Talk to your care team if you have: • A nosebleed lasting more than 5 minutes despite pressure • A cut that continues to bleed • Heavy gum bleeding when brushing or flossing • Sudden or severe headache • Blood in your urine or stool • Blood in your spit after coughing
Low Red Blood Cell (RBC) Count and Hemoglobin (Hgb) (Anemia)	Description: Red blood cells carry oxygen throughout your body. Low levels can make you feel tired, weak, dizzy, or short of breath. Recommendations: • Aim for 7 to 8 hours of sleep each night. • Do not drive, operate heavy machinery, or do other dangerous activities if you are very tired. • Balance activity and rest — stay as active as you can, but rest when needed. • Eat a balanced diet and follow any nutrition or supplement advice from your care team. Talk to your care team if you have: • Shortness of breath • Dizziness or fainting • Fast or irregular heartbeats • Sudden or severe headache

Brought to you by: ACCC Association of Cancer Care Centers, HOPA Hematology/Oncology Pharmacy Association, NCODA Network for Collaborative Oncology Development & Advancement, ONS Oncology Nursing Society

<table>
<tr><td>Fatigue</td><td>Description: Fatigue is a constant and sometimes strong feeling of tiredness.

Recommendations:
Routine exercise can help reduce fatigue. Talk with your care team to find the right type and amount of activity for you.
Ask family and friends for help with daily tasks and for emotional support.
Try healthy ways to feel better, such as meditation, journaling, yoga, or guided imagery, to reduce anxiety and improve well-being.
Aim for 7 to 8 hours of sleep each night. Limit daytime naps to help you sleep better at night.
Do not drive, operate heavy machinery, or do other potentially dangerous activities if you are very tired.
Talk to your care team if you have:
Tiredness that affects your daily life or prevents you from doing normal activities
Tiredness that does not get better with rest
Dizziness or weakness along with severe tiredness</td></tr>
<tr><td>Nausea and Vomiting</td><td>Description: Nausea is an uncomfortable feeling in your stomach or the need to throw up. You may or may not vomit.

Recommendations:
Eat smaller, more frequent meals.
Avoid fatty, fried, spicy, or highly sweet foods.
Eat bland foods at room temperature and drink clear liquids.
If you vomit, start with small sips of water, broth, or other clear liquids. If these stay down, try soft foods (such as gelatin, plain cornstarch pudding, yogurt, strained soup, or strained cooked cereal) and gradually return to solid foods.
Your care team may prescribe medicine for these symptoms.
Talk to your care team if you have:
Vomiting for more than 24 hours
Nonstop vomiting
Signs of dehydration (very thirsty, dry mouth, dizziness, or dark urine)
Blood in your vomit or vomit that looks like coffee grounds
Severe stomach pain that does not go away after vomiting</td></tr>
</table>

Brought to you by:
ACCC Association of Cancer Care Centers
HOPA Hematology/Oncology Pharmacy Association
NCODA Network for Collaborative Oncology Development & Advancement
ONS Oncology Nursing Society

Low Appetite and Weight Loss	Description: Loss of appetite can lead to weight loss and low energy. Small changes in when and what you eat can help maintain strength and nutrition. Recommendations: • Be as active as you can. Do light physical activity before a meal (check with your care team before starting an exercise program). • Note times of day when your appetite is best and eat your largest meal then. • Eat 5 or 6 small meals or snacks each day. • Choose high-protein foods (beans, chicken, fish, meat, yogurt, tofu, eggs). Eat protein first during meals. • Choose higher-calorie foods (avoid “low-fat”, “fat-free”, or “diet” options when trying to gain/maintain weight). • If you feel full quickly, avoid drinking 30 minutes before a meal and drink liquids between meals; choose calorie-containing drinks rather than diet drinks. • Have a bedtime snack that’s easy to digest (for example, peanut butter and crackers). If you have reflux, wait at least 1 hour before lying down. • Try nutritious beverages (high-protein shakes or smoothies) if solid food is unappealing. • Ask your care team about liquid nutrition supplements and ways to add protein or calories (protein powder, yogurt, ice cream). Talk to your care team if you have: • Unintentional weight loss • Little or no appetite for several days • Excessive tiredness or low energy
Liver Problems	Description: Treatment can cause liver injury. Your care team may check your liver with blood tests before and during treatment. Talk to your care team if you have: • Yellowing of your skin or the white part of your eyes (jaundice) • Severe nausea or vomiting • Pain on the right side of your stomach area (abdomen) • Dark, tea-colored urine • Bleeding or bruising more easily than normal

Brought to you by: ACCC Association of Cancer Care Centers™ HOPA Hematology/Oncology Pharmacy Association NCODA Network for Collaborative Oncology Development & Advancement ONS Oncology Nursing Society

Diarrhea	Description: Diarrhea means having bowel movements more often than usual. Your stools may also be loose or watery. If you have diarrhea: • Keep track of how often you have diarrhea. • Drink 8 to 10 glasses of water or fluids each day. Do this unless your care team tells you otherwise. • Eat small meals with low-fiber foods. Try bananas, applesauce, potatoes, chicken, rice, or toast. • Avoid high-fiber foods. These include raw vegetables, fruit, and whole grains. • Avoid foods that may cause gas. These include broccoli and beans. • Avoid milk, yogurt, and other dairy foods. • Avoid spicy, fried, or greasy foods. • Your care team may recommend medicine for diarrhea. One example is loperamide (Imodium). Talk to your care team if you have: • An increase of 4 or more bowel movements than usual in one day • Dizziness or lightheadedness while you have diarrhea • Signs of dehydration, such as feeling very thirsty, dizzy, or having dark urine • Blood in your stool
Kidney Problems	Description: Treatment can cause kidney problems, including damage to the kidneys and decreased kidney function. Your care team will monitor your kidney function during treatment. Recommendations: • Drink 8 to 10 glasses of water or other fluids each day, unless your care team tells you otherwise. • Your care team may give you fluids and electrolytes with your treatment. Talk to your care team if you have: • Decrease in your amount of urine • Blood in your urine • Swelling of your ankles • Loss of appetite

Brought to you by:
ACCC Association of Cancer Care Centers
HOPA Hematology/Oncology Pharmacy Association
NCODA Network for Collaborative Oncology Development & Advancement
ONS Oncology Nursing Society

High Blood Sugar (Hyperglycemia)	Description: Treatment may cause high blood sugar levels. Your care team may perform blood tests to check for these changes and treat you if needed. Recommendations: • Eat a well-balanced diet. • Limit sugary drinks and foods. • Eat smaller, more frequent meals. • Be physically active for at least 30 minutes most days. • Your care team may ask you to check your blood sugar at home. If you are already doing this, they may ask you to do it more frequently. Talk to your care team if you have: • Frequent urination • Drowsiness • Increased thirst • Loss of appetite • Blurred vision • Fruity smell on your breath • Confusion • Nausea, vomiting, or stomach pain • It becomes harder to control your blood sugar
Changes in Electrolytes and Other Laboratory Results	Description: Treatment may cause low levels of phosphate, sodium, and calcium in your blood. Your care team will perform blood tests to check for these changes and will treat you if needed. Talk to your care team if you have: • New or worsening muscle weakness, cramps, or bone pain • Trouble breathing or shortness of breath • Slurred speech, or severe dizziness • Severe fatigue or inability to perform daily activities • New or worsening confusion, drowsiness, or trouble thinking • Severe headache or nausea/vomiting • Seizures or fainting • Sudden or new weakness, unsteadiness, or trouble walking • Severe muscle cramps or spasms • Tingling or numbness around the mouth, hands, or feet • Fast, irregular, or pounding heartbeat • Sudden weight gain • Swelling of your arms, hands, legs, or ankles

Brought to you by:
ACCC Association of Cancer Care Centers
HOPA Hematology/Oncology Pharmacy Association
NCODA Network for Collaborative Oncology Development & Advancement
ONS Oncology Nursing Society

Select Rare Side Effects

Side Effect	Talk to Your Care Team if You Have Any of These Signs or Symptoms
Eye Problems (Cataracts)	Treatment may cause new cataracts or make existing cataracts worse. Cataracts cause the lens of your eye to become cloudy, making it harder to see clearly. • Blurred or cloudy vision • Double vision • Sensitivity to light or glare • Vision changes
Brain Problems	• Headache • Dizziness • Confusion • Memory problems • Difficulty concentrating • Fatigue • Seizures • Muscle weakness • Tremors • Changes in mood or behavior • Sensory disturbances (such as tingling or numbness)

Before starting treatment, ask your care team when to call 9-1-1 or seek emergency help.
If you experience any new, worsening, or uncontrolled side effects, contact your care team immediately.

Intimacy, Fertility, Pregnancy, and Breastfeeding

- Treatment may **change how you feel about intimacy and your body.** However, physical closeness—such as holding hands and hugging—remains safe. It is common to have questions about intimacy. If needed, talk to your care team for guidance.
- Treatment can affect your **ability to have children**. It may damage your reproductive organs or stop them from working. If you are worried about fertility, talk to your care team before starting treatment.
- Treatment may **harm an unborn baby**.
    - If you are able to become pregnant:
        - Your care team will verify your pregnancy status before treatment.
        - Use an effective method of birth control during treatment and for 1 week after your last dose of selinexor.
        - If you think you might be pregnant or if you become pregnant, tell your care team right away.
    - If your partner is able to become pregnant, use an effective method of birth control—such as condoms—during treatment and for 1 week after your last dose of selinexor.
- **Do not breastfeed** during treatment and for 1 week after your last dose of selinexor.

Brought to you by: ACCC Association of Cancer Care Centers, HOPA Hematology/Oncology Pharmacy Association, NCODA Network for Collaborative Oncology Development & Advancement, ONS Oncology Nursing Society

Additional Information

- **Tell your care team about all the medicines you take**.
  This includes prescriptions, over-the-counter drugs, vitamins, and herbal products. Before starting any new medicine, supplement, or vaccine, ask your care team first.
- **This Patient Education Sheet may not describe all possible side effects.**
  Call your care team for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Notes

Updated Date: October 29, 2026

Scan the QR code below to access this education sheet.

Important Notice:
The Association of Cancer Care Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), Network for Collaborative Oncology Development & Advancement, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated to gather information and develop this patient education guide. This guide is a summary of the medication, based on information provided by the drug manufacturer and other sources.

This guide does not cover all information available on the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this medication. It should not be used as a substitute for the advice of a qualified healthcare professional. The provision of this guide is for informational purposes only. It does not constitute or imply endorsement, recommendation, or favoring of this medication by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. All decisions regarding this medication should be made with the guidance and direction of a qualified healthcare professional.

Permission:
Patient Education Sheets are provided as a free educational resource for patients with cancer and their caregivers in need of concise, easy-to-understand information about cancer therapy. Healthcare providers may copy and distribute the sheets to patients and direct them to the Patient Education Sheets website. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holders. Permission requests, including direct linking from Electronic Health Records, and licensing inquiries should be emailed to patienteducationsheets@ncoda.org.

Copyright © 2026 by Network for Collaborative Oncology Development & Advancement, Inc. All rights reserved.

PES-226

Brought to you by:
ACCC Association of Cancer Care Centers
HOPA Hematology/Oncology Pharmacy Association
NCODA Network for Collaborative Oncology Development & Advancement
ONS Oncology Nursing Society