

Vismodegib

Care Team Contact Information: _____

Pharmacy Contact Information: _____

Diagnosis: _____

- This treatment is often used for a type of skin cancer called basal cell carcinoma (BCC).
- It may also be used for other reasons.

Goal of Treatment: _____

- Treatment may continue for a certain time period, until it no longer works, or until side effects are no longer controlled.

Treatment Regimen

Treatment Name	How the Treatment Works	How the Treatment is Given
Vismodegib (VIS-moh-DEH-gib): Erivedge (AYR-ih-vej)	Slows down or stops the growth of cancer cells by blocking a specific protein that helps them survive.	Capsule taken by mouth.

Treatment Administration and Schedule

Your vismodegib dosing instructions:

- Vismodegib comes in 1 capsule strength: 150 mg.
- Your dose might differ, but vismodegib is typically taken as one 150 mg capsule once daily.
- Take vismodegib 1 time each day, around the same time each day.
- You can take vismodegib with or without food.
- Swallow vismodegib capsules whole. Do not open or crush the capsules.
- Do not change your dose or stop taking vismodegib unless your care team tells you to.
- Your care team may tell you to temporarily stop or completely stop taking vismodegib if you develop certain side effects.
- If you miss a dose, skip the missed dose. Just take your next scheduled dose.
- If you vomit after taking a dose of vismodegib, do not make up the dose. Take your next dose at your regularly scheduled time.
- If you take too much vismodegib, call your care team or go to the nearest hospital emergency room right away.

Storage and Handling of vismodegib:

- Store vismodegib at room temperature between 68°F and 77°F (20°C and 25°C).
- Keep the bottle tightly closed to protect vismodegib from moisture.
- People who are or may be pregnant should wear gloves when handling vismodegib.
- Keep vismodegib and all medicines out of the reach of children and pets.
- Ask your care team how to safely throw away any unused vismodegib.

Appointments: Appointments may include regular check-ups with your care team, lab visits, and imaging tests. It's important to keep your appointments whenever you can. If you miss any appointments, call your care provider as soon as possible to reschedule your appointment.

Supportive Care to Prevent and Treat Side Effects

Description	Supportive Care Taken at Home
To help prevent or treat nausea and vomiting	<hr/> <hr/> <hr/>
Other	<hr/> <hr/> <hr/>

Common Side Effects

Side Effect	Important Information
Fatigue	<p>Description: Fatigue is a constant and sometimes strong feeling of tiredness.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Routine exercise can help reduce fatigue. Talk with your care team to find the right type and amount of activity for you. • Ask family and friends for help with daily tasks and for emotional support. • Try healthy ways to feel better, such as meditation, journaling, yoga, or guided imagery, to reduce anxiety and improve well-being. • Aim for 7 to 8 hours of sleep each night. Limit daytime naps to help you sleep better at night. • Do not drive, operate heavy machinery, or do other potentially dangerous activities if you are very tired. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • Tiredness that affects your daily life or prevents you from doing normal activities • Tiredness that does not get better with rest • Dizziness or weakness along with severe tiredness
Nausea and Vomiting	<p>Description: Nausea is an uncomfortable feeling in your stomach or the need to throw up. You may or may not vomit.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Eat smaller, more frequent meals. • Avoid fatty, fried, spicy, or highly sweet foods. • Eat bland foods at room temperature and drink clear liquids. • If you vomit, start with small sips of water, broth, or other clear liquids. If these stay down, try soft foods (such as gelatin, plain cornstarch pudding, yogurt, strained soup, or strained cooked cereal) and gradually return to solid foods. • Your care team may prescribe medicine for these symptoms. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • Vomiting for more than 24 hours • Nonstop vomiting • Signs of dehydration (very thirsty, dry mouth, dizziness, or dark urine) • Blood or coffee-ground-like appearance in your vomit • Severe stomach pain that does not go away after vomiting

<p>Diarrhea</p>	<p>Description: Diarrhea is loose, watery stools or more frequent bowel movements than usual. It can cause dehydration and weakness.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Keep track of how often you go to the bathroom each day. • Drink 8 to 10 glasses of water or other fluids daily, unless your care team tells you otherwise. • Eat small meals of mild, low-fiber foods (such as bananas, applesauce, potatoes, chicken, rice, and toast). • If you have diarrhea, avoid high-fiber foods (such as raw vegetables, fruits, and whole grains), gas-producing foods (such as broccoli and beans), dairy (such as milk and yogurt), and spicy, fried, or greasy foods. • Your care team may recommend an antidiarrheal medicine such as loperamide (Imodium). <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • 4 or more bowel movements than normal in 24 hours • Dizziness or lightheadedness while having diarrhea • Signs of dehydration (very thirsty, dry mouth, dizziness, or dark urine) • Bloody diarrhea
<p>Weight Loss</p>	<p>Description: Loss of appetite can lead to weight loss and low energy. Small changes in when and what you eat can help maintain strength and nutrition.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Be as active as you can. Do light physical activity before a meal (check with your care team before starting an exercise program). • Note times of day when your appetite is best and eat your largest meal then. • Eat 5–6 small meals or snacks each day. • Choose high-protein foods (beans, chicken, fish, meat, yogurt, tofu, eggs). Eat protein first during meals. • Choose higher-calorie foods (avoid “low-fat,” “fat-free,” or “diet” options when trying to gain/maintain weight). • If you feel full quickly, avoid drinking 30 minutes before a meal and drink liquids between meals; choose calorie-containing drinks rather than diet drinks. • Have a bedtime snack that’s easy to digest (for example, peanut butter and crackers). If you have reflux, wait at least 1 hour before lying down. • Try nutritious beverages (high-protein shakes or smoothies) if solid food is unappealing. • Ask your care team about liquid nutrition supplements and ways to add protein or calories (protein powder, yogurt, ice cream). <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • Unintentional weight loss • Little or no appetite for several days • Excessive tiredness or low energy

<p>Taste Changes</p>	<p>Description: Taste changes, such as loss of taste or taste sensitivity, may occur. Foods that used to taste good may no longer be appealing or certain tastes, such as metallic tastes, may become more noticeable.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Choose appealing foods based on appearance and smell. • Use plastic utensils if food tastes metallic. • Add spices or juices to enhance food flavors. • Suck on mints or chew gum to improve taste. • Brush your teeth before and after eating with a soft toothbrush. • Avoid smoking. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • Trouble eating • Weight loss
<p>Muscle Spasm or Twitching</p>	<p>Description: Treatment can sometimes make your muscles twitch or cramp up. This might feel like a little jump or twitch in a part of your body that you can't control, or a feeling of tightness that can be painful.</p> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • Muscle cramps and stiffness • Muscle pain
<p>Hair Loss (Alopecia)</p>	<p>Description: Hair loss or thinning may begin days to weeks after treatment starts and usually grows back later. New growth can be a different color or texture and may not look the same as before.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Consider a short haircut before treatment and use scarves, hats, or wigs for comfort and confidence. • Keep your head covered outdoors to protect it from the sun and cold; use sunscreen on your uncovered scalp. • Use gentle haircare: mild shampoo, soft brush, and avoid heat styling and harsh treatments. • Ask your care team about wig prescriptions or resources for head coverings. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • No hair regrowth months after treatment ends • Concern about hair changes or need help finding a wig or support resources

Select Rare Side Effects

Side Effect	Talk to Your Care Team if You Have Any of These Signs or Symptoms
Severe Skin Reactions	<ul style="list-style-type: none"> • A red, itchy, skin rash • Peeling of your skin or blisters • Severe itching • Fever of 100.4°F (38°C) or higher

Before starting treatment, ask your care team when to call 9-1-1 or seek emergency help. If you experience any new, worsening, or uncontrolled side effects, contact your care team immediately.

Intimacy, Fertility, Pregnancy, and Breastfeeding

- Treatment may **change how you feel about intimacy and your body**. However, physical closeness—such as holding hands and hugging—remains safe. It is common to have questions about intimacy. If needed, talk to your care team for guidance.
- Treatment may affect your **ability to have children**. It may damage your reproductive organs or stop them from working. If you are worried about fertility, talk to your care team before starting treatment.
- Treatment may **harm an unborn baby**.
 - If you are able to become pregnant, take a pregnancy test before starting treatment.
 - Use an effective method of birth control during treatment and for 24 months after your last dose of vismodegib.
 - If you think you might be pregnant or if you become pregnant, tell your care team right away.
 - If your partner is able to become pregnant, use an effective method of birth control—such as condoms—during treatment and 3 months after your last dose of vismodegib.
 - Do not donate semen during treatment with vismodegib and for 3 months after your final dose.
- **Do NOT breastfeed** during treatment and for 24 months after your last dose of vismodegib.

Additional Information

- **Tell your care team about all the medicines you take.**
This includes prescriptions, over-the-counter drugs, vitamins, and herbal products. Before starting any new medicine, supplement, or vaccine, ask your care team first.
- **Do not donate blood or blood products** while receiving vismodegib and for 24 months after the final dose of vismodegib.
- **This Patient Education Sheet may not describe all possible side effects.**
Call your care team for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Notes

Updated Date: May 19, 2026

Scan the QR code below to access this education sheet.



Important notice: The Association of Cancer Care Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), Network for Collaborative Oncology Development & Advancement, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated in gathering information for and developing this patient education guide. This guide represents a brief summary of the medication derived from information provided by the drug manufacturer and other resources.

This guide does not cover all existing information related to the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this medication and should not substitute for the advice of a qualified healthcare professional. Provision of this guide is for informational purposes only and does not constitute or imply endorsement, recommendation, or favoring of this medication by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional.

Permission: Patient Education Sheets are provided as a free educational resource for patients with cancer and their caregivers in need of concise, easy to understand information about cancer therapy. Healthcare providers are permitted to copy and distribute the sheets to patients as well as direct patients to the Patient Education Sheets website. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holders. Permission requests, including direct linking from Electronic Health Records, and licensing inquiries should be emailed to patienteducation sheets@ncoda.org.

Copyright © 2026 by Network for Collaborative Oncology Development & Advancement, Inc. All rights reserved.

PES-276