

Glofitamab

Care Team Contact Information: _____

Pharmacy Contact Information: _____

Diagnosis: _____

- This treatment is used for diffuse large B-cell lymphoma (DLBCL) or large B-cell lymphoma (LBCL).
- It may also be used for other reasons.

Goal of Treatment: _____

- Treatment may continue for a certain time period, until it no longer works, or until side effects are no longer controlled.

Treatment Regimen

Treatment Name	How the Treatment Works	How the Treatment is Given
Glofitamab ((gloh-FIH-tah-mab): Columvi (koh-LOOM-vee)	Binds immune cells (T-cells) to cancer cells so T-cells can more effectively attack and destroy them.	Infusion into a vein (intravenous (IV) infusion).

Treatment Administration and Schedule

Treatment is typically repeated every 3 weeks. This length of time is called a "cycle."

Due to the risk of cytokine release syndrome (CRS), you will receive glofitamab on a **"step-up dosing schedule."**

- A single dose of a medicine called obinutuzumab will be given to you on the first day of your first treatment cycle (Day 1 of Cycle 1).
- You will start the glofitamab step-up dosing schedule a week after the obinutuzumab dose.
 - The step-up dosing schedule is when you receive smaller "step-up" doses of glofitamab on Day 8 and Day 15 of Cycle 1. This is to help reduce your risk of CRS.
 - If your dose is delayed for any reason, you may need to repeat the "step-up dosing schedule."
- You may be hospitalized during your infusion and for 24 hours after receiving the first step-up dose on Day 8.
- You may be hospitalized during your infusion and for 24 hours after receiving the second step-up dose on Day 15 if you experienced CRS during the first step-up dose.
- You will receive your first full dose of glofitamab a week after the second step-up dose (this will be Day 1 of Cycle 2).

Treatment Administration and Schedule (Continued)

Cycle 1

Treatment Name	Cycle 1							
	Day 1	Day 2	...	Day 8	...	Day 15	...	Day 21
Obinutuzumab	✓							
Glofitamab				✓ (Step-Up Dose 1)		✓ (Step-Up Dose 2)		

Cycles 2 to 12

Treatment Name	Cycle 2								Next Cycle
	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	...	Day 21	Day 1
Glofitamab	✓ (First Full Dose)								✓

Appointments

Appointments may include regular check-ups with your care team, treatment appointments, lab visits, and imaging tests. It's important to keep your appointments whenever you can. If you miss an appointment, call your care team as soon as possible to reschedule it.

Supportive Care to Prevent and Treat Side Effects

Description	Supportive Care Given at the Clinic or Hospital	Supportive Care Taken at Home
To help lower the risk of Cytokine Release Syndrome (CRS)		
To help lower the risk of infections		
To help lower the risk of Tumor Lysis Syndrome (TLS)		
Other		

Common Side Effects

Side Effect	Important Information
Cytokine Release Syndrome (CRS) (Boxed Warning)	Description: CRS happens when your immune system becomes very active after treatment. Most CRS events are mild, get better with treatment, and happen during the first few doses. However, some CRS events can be serious and life-threatening. Symptoms can include fever, chills, fatigue, headache, dizziness or feeling lightheaded, or difficulty breathing. Recommendations: - Keep a symptom diary to record any new or worsening symptoms such as fever, chills, fatigue, or difficulty breathing. - Your care team may ask you to monitor your temperature, blood pressure, heart rate, or oxygen level. - Stay hydrated by drinking plenty of fluids to help manage symptoms and support overall health. - If you develop CRS, do not drive or operate dangerous machinery until your care team says it is safe. - Your care team may prescribe medications for CRS. Talk to your care team if you have: - Fever of 100.4°F (38°C) or higher - Trouble breathing - Chills - Dizziness or light-headedness - Fast heartbeat - Headache Note: Your care team may have specific numbers for blood pressure, heart rate, and blood oxygen levels. If your numbers go beyond those limits, call your care team or get emergency help.
Low White Blood Cell (WBC) Count (Neutropenia) and Increased Risk of Infection	Description: WBCs help protect your body from infections. A low WBC count increases your risk of infection. Recommendations: - Wash your hands often and bathe regularly. - Avoid crowded places and close contact with people who are sick. - Follow food safety and wound care advice from your care team. - Your care team may prescribe medicine to help your WBCs recover. Talk to your care team if you have: - Fever of 100.4°F (38°C) or higher - Chills - New or worsening cough or sore throat - Painful urination or signs of a urinary infection - Feeling much more tired than usual - Red, swollen, warm, or painful areas on the skin (possible skin infection)

Low Platelet Count (Thrombocytopenia)	Description: Platelets help your blood clot and wounds heal. A low platelet count increases your risk of bruising and bleeding. Recommendations: • Blow your nose gently and avoid picking it. • Brush your teeth gently with a soft toothbrush and keep good oral hygiene. • Use an electric razor for shaving and a nail file instead of nail clippers. • Avoid over-the-counter medicines that can increase bleeding risk (for example, nonsteroidal anti-inflammatory drugs (NSAIDs) like ibuprofen). • Talk with your care team or dentist before medical or dental procedures — you may need to pause treatment. Talk to your care team if you have: • A nosebleed lasting more than 5 minutes despite pressure • A cut that continues to bleed • Heavy gum bleeding when brushing or flossing • Sudden or severe headache • Blood in your urine or stool • Blood in your spit after coughing
Low Red Blood Cell (RBC) Count and Hemoglobin (Hgb) (Anemia)	Description: RBCs and Hgb carry oxygen to your body's tissues and remove carbon dioxide. Low RBC or Hgb (anemia) can make you feel weak or very tired, or may make you look pale. Recommendations: • Aim for 7 to 8 hours of sleep each night. • Do not drive, operate heavy machinery, or do other dangerous activities if you are very tired. • Balance activity and rest — stay as active as you can, but rest when needed. • Eat a balanced diet and follow any nutrition or supplement advice from your care team. Talk to your care team if you have: • Shortness of breath • Dizziness or fainting • Fast or irregular heartbeats • Sudden or severe headache

Fatigue	Description: Fatigue is a constant and sometimes strong feeling of tiredness. Recommendations: • Routine exercise can help reduce fatigue. Talk with your care team to find the right type and amount of activity for you. • Ask family and friends for help with daily tasks and for emotional support. • Try healthy ways to feel better, such as meditation, journaling, yoga, or guided imagery, to reduce anxiety and improve well-being. • Aim for 7 to 8 hours of sleep each night. Limit daytime naps to help you sleep better at night. • Do not drive, operate heavy machinery, or do other potentially dangerous activities if you are very tired. Talk to your care team if you have: • Tiredness that affects your daily life or prevents you from doing normal activities • Tiredness that does not get better with rest • Dizziness or weakness along with severe tiredness
Muscle or Joint Pain or Weakness	Description: Muscle pain is soreness, aching, cramps, stiffness, tenderness, or weakness in one or more muscles. Joint pain is pain, stiffness, swelling, or reduced movement where two bones meet. Recommendations: • Keep a pain diary: note pain levels, locations, and activities that make it better or worse. • Do gentle exercise (walking, stretching, yoga) to maintain mobility and strength. Check with your care team before starting a new activity. • Use a warm compress for stiff muscles or a cold pack to reduce swelling and numb pain—use what helps the area. • Your care team may recommend or prescribe medicines, including over-the-counter pain relievers. Talk to your care team if you have: • Pain you cannot control with usual measures • Swelling, redness, or warmth in a joint • New weakness • Trouble walking or moving

Select Rare Side Effects

Side Effect	Talk to Your Care Team if You Have Any of These Signs or Symptoms
Neurologic Problems (Boxed Warning)	Treatment can cause serious neurologic (brain and nerve) problems that can be life-threatening and may lead to death. These problems can include a condition called Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS). Neurologic problems may occur days or weeks after treatment. Your care team may refer you to a specialist who treats neurologic problems. If you develop neurologic side effects, do not drive or operate dangerous machinery until your care team says it is safe. • Headache • Agitation, trouble staying awake, confusion or disorientation, seeing or hearing things that are not real (hallucinations) • Trouble speaking, writing, thinking, remembering things, paying attention, or understanding things • Problems walking, muscle weakness, shaking (tremors), loss of balance, or muscle spasms • Numbness and tingling (feeling like "pins and needles") • Burning, throbbing, or stabbing pain • Changes in your handwriting • Seizures
Growth in Your Tumor or Worsening of Tumor-Related Problems (Tumor Flare)	• Tender or swollen lymph nodes • Pain or swelling at the site of the tumor • Chest pain • Cough • Trouble breathing

Before starting treatment, ask your care team when to call 9-1-1 or seek emergency help.
If you experience any new, worsening, or uncontrolled side effects, contact your care team immediately.

Intimacy, Pregnancy, and Breastfeeding

- Treatment may **change how you feel about intimacy and your body**. However, physical closeness—such as holding hands and hugging—remains safe. It is common to have questions about intimacy. If needed, talk to your care team for guidance.
- Treatment may **harm an unborn baby**.
 - If you are able to become pregnant:
 - Your care team will verify your pregnancy status before treatment.
 - Use an effective method of birth control during treatment and for 1 month after your last dose of glofitamab.
 - If you think you might be pregnant or if you become pregnant, tell your care team right away.
 - If your partner is able to become pregnant, use an effective method of birth control—such as condoms—during treatment with glofitamab.
- **Do not breastfeed** during treatment and for 1 month after your last dose of glofitamab.

Additional Information

- **Tell your care team about all the medicines you take.**
This includes prescriptions, over-the-counter drugs, vitamins, and herbal products. Before starting any new medicine, supplement, or vaccine, ask your care team first.
- **Always carry the glofitamab Patient Wallet Card** with you and show it to all of your healthcare providers.
The glofitamab Patient Wallet Card lists the signs and symptoms of CRS that you should get medical help for right away.
- **This Patient Education Sheet may not describe all possible side effects.**
Call your care team for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Notes

Updated Date: August 17, 2026

Scan the QR code below to access this education sheet.

**Important Notice:**

The Association of Cancer Care Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), Network for Collaborative Oncology Development & Advancement, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated to gather information and develop this patient education guide. This guide is a summary of the medication, based on information provided by the drug manufacturer and other sources.

This guide does not cover all information available on the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this medication. It should not be used as a substitute for the advice of a qualified healthcare professional. The provision of this guide is for informational purposes only. It does not constitute or imply endorsement, recommendation, or favoring of this medication by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. All decisions regarding this medication should be made with the guidance and direction of a qualified healthcare professional.

Permission:

Patient Education Sheets are provided as a free educational resource for patients with cancer and their caregivers in need of concise, easy-to-understand information about cancer therapy. Healthcare providers may copy and distribute the sheets to patients and direct them to the Patient Education Sheets website. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holders. Permission requests, including direct linking from Electronic Health Records, and licensing inquiries should be emailed to patienteducationsheets@ncoda.org.

Copyright © 2026 by Network for Collaborative Oncology Development & Advancement, Inc. All rights reserved.

PES-106