

Ribociclib

Care Team Contact Information: _____

Pharmacy Contact Information: _____

Diagnosis: _____

- This treatment is often used for certain types of breast cancer.
- It may also be used for other reasons.

Goal of Treatment: _____

- Treatment may continue for a certain time period, until it no longer works, or until side effects are no longer controlled.

Treatment Regimen

Treatment Name	How the Treatment Works	How the Treatment is Given
Ribociclib (RY-boh-SY-klib): Kisqali (kis-KAH-lee)	Slows down or stops the growth of cancer cells by blocking a specific protein that helps them survive.	Tablet(s) taken by mouth.

Treatment Administration and Schedule

Ribociclib is typically taken by mouth 1 time a day for 21 days, followed by a 7-day rest period. This 28-day length of time is called a “cycle”.

Your ribociclib dosing instructions:

- Ribociclib comes in 1 tablet strength: 200 mg. There are 3 different sized tablet therapy packs: 200 mg, 400 mg, and 600 mg. Your care team will tell you which therapy pack to take.
- Your dose might differ, but ribociclib is typically taken as
 - 400 mg (two 200 mg tablets) once daily on days 1 to 21 of a 28-day treatment cycle OR
 - 600 mg (three 200 mg tablets) once daily on days 1 to 21 of a 28-day treatment cycle
- Take ribociclib 1 time each day, around the same time each day, preferably in the morning.
- Take ribociclib with or without food.
- Swallow ribociclib tablets whole. Do not chew, crush, or split ribociclib tablets before swallowing them.
- Do not take any ribociclib tablets that are broken, cracked, or that look damaged.
- Do not change your dose or stop taking ribociclib unless your care team tells you to.

Treatment Administration and Schedule Continued

- Your care team may tell you to change your dose, temporarily stop, or completely stop taking ribociclib if you develop certain side effects.
- If you miss a dose of ribociclib or vomit after taking a dose of ribociclib, do not take another dose on that day. Take your next dose at your regular time.
- If you take too much ribociclib, call your care team or go to the nearest hospital emergency room right away.

Treatment Name	Cycle 1, Days																						Next Cycle
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22-28	Day 1
Ribociclib	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Week-long Break	✓

Storage and Handling of Ribociclib

- Store ribociclib at room temperature between 68°F and 77°F (20°C and 25°C).
- Store ribociclib in the original blister package in order to protect from moisture.
- Keep ribociclib and all medicines out of the reach of children and pets.
- Ask your care team how to safely throw away any unused ribociclib.

Appointments

Appointments may include regular check-ups with your care team, lab visits, and imaging tests. It's important to keep your appointments whenever you can. If you miss any appointments, call your care provider as soon as possible to reschedule your appointment.

Supportive Care to Prevent and Treat Side Effects

Description	Supportive Care Taken at Home
To help prevent or treat nausea and vomiting	
Other	

Common Side Effects

Side Effect	Important Information
Low White Blood Cell (WBC) Count (Neutropenia) and Increased Risk of Infection	Description: WBCs help protect your body from infections. A low WBC count increases your risk of getting infections. Recommendations: - Wash your hands often and bathe regularly. - Avoid crowded places and close contact with people who are sick. - Follow food safety and wound care advice from your care team. - Your care team may prescribe medicine to help your WBCs recover. Talk to your care team if you have: - Fever of 100.4°F (38°C) or higher - Chills - New or worsening cough or sore throat - Painful urination or signs of a urinary infection - Feeling much more tired than usual - Red, swollen, warm, or painful areas on the skin (possible skin infection)
Low Platelet Count (Thrombocytopenia)	Description: Platelets help your blood clot and wounds heal. A low platelet count increases your risk of bruising and bleeding. Recommendations: - Blow your nose gently and avoid picking it. - Brush your teeth gently with a soft toothbrush and keep good oral hygiene. - Use an electric razor for shaving and a nail file instead of nail clippers. - Avoid over-the-counter medicines that can increase bleeding risk (for example, nonsteroidal anti-inflammatory drugs (NSAIDs) like ibuprofen). - Talk with your care team or dentist before medical or dental procedures — you may need to pause treatment. Talk to your care team if you have: - A nosebleed lasting more than 5 minutes despite pressure - A cut that continues to bleed - Heavy gum bleeding when brushing or flossing - Sudden or severe headache - Blood in your urine or stool - Blood in your spit after coughing
Low Red Blood Cell (RBC) Count and Hemoglobin (Hgb) (Anemia)	Description: RBCs and Hgb carry oxygen to your body's tissues and remove carbon dioxide. Low RBC or Hgb (anemia) can make you feel weak, very tired, or look pale. Recommendations: - Aim for 7 to 8 hours of sleep each night. - Do not drive, operate heavy machinery, or do other dangerous activities if you are very tired. - Balance activity and rest — stay as active as you can, but rest when needed. - Eat a balanced diet and follow any nutrition or supplement advice from your care team. Talk to your care team if you have: - Shortness of breath - Dizziness or fainting - Fast or irregular heartbeats - Sudden or severe headache

Fatigue	Description: Fatigue is a constant and sometimes strong feeling of tiredness. Recommendations: • Routine exercise can help reduce fatigue. Talk with your care team to find the right type and amount of activity for you. • Ask family and friends for help with daily tasks and for emotional support. • Try healthy ways to feel better, such as meditation, journaling, yoga, or guided imagery, to reduce anxiety and improve well-being. • Aim for 7 to 8 hours of sleep each night. Limit daytime naps to help you sleep better at night. • Do not drive, operate heavy machinery, or do other potentially dangerous activities if you are very tired. Talk to your care team if you have: • Tiredness that affects your daily life or prevents you from doing normal activities • Tiredness that does not get better with rest • Dizziness or weakness along with severe tiredness
Nausea and Vomiting	Description: Nausea is an uncomfortable feeling in your stomach or the need to throw up. You may or may not vomit. Recommendations: • Eat smaller, more frequent meals. • Avoid fatty, fried, spicy, or highly sweet foods. • Eat bland foods at room temperature and drink clear liquids. • If you vomit, start with small sips of water, broth, or other clear liquids. If these stay down, try soft foods (such as gelatin, plain cornstarch pudding, yogurt, strained soup, or strained cooked cereal) and gradually return to solid foods. • Your care team may prescribe medicine for these symptoms. Talk to your care team if you have: • Vomiting for more than 24 hours • Nonstop vomiting • Signs of dehydration (very thirsty, dry mouth, dizziness, or dark urine) • Blood or coffee-ground-like appearance in your vomit • Severe stomach pain that does not go away after vomiting
Liver Problems	Description: Treatment can cause liver injury. Your care team may check your liver with blood tests before and during treatment. Talk to your care team if you have: • Yellowing of your skin or the white part of your eyes (jaundice) • Severe nausea or vomiting • Pain on the right side of your stomach area (abdomen) • Dark, tea-colored urine • Bleeding or bruising more easily than normal

Diarrhea	Description: Diarrhea is loose, watery stools or more frequent bowel movements than usual. It can cause dehydration and weakness. Recommendations: • Keep track of how often you go to the bathroom each day. • Drink 8 to 10 glasses of water or other fluids daily, unless your care team tells you otherwise. • Eat small meals of mild, low-fiber foods (such as bananas, applesauce, potatoes, chicken, rice, and toast). • If you have diarrhea, avoid high-fiber foods (such as raw vegetables, fruits, and whole grains), gas-producing foods (such as broccoli and beans), dairy (such as milk and yogurt), and spicy, fried, or greasy foods. • Your care team may recommend an antidiarrheal medicine such as loperamide (Imodium). Talk to your care team if you have: • 4 or more bowel movements than normal in 24 hours • Dizziness or lightheadedness while having diarrhea • Signs of dehydration (very thirsty, dry mouth, dizziness, or dark urine) • Bloody diarrhea
Kidney Problems	Description: Treatment can cause kidney problems, including damage to the kidneys and decreased kidney function. Your care team will monitor your kidney function during treatment. Recommendations: • Drink 8 to 10 glasses of water or other fluids each day, unless your care team tells you otherwise. • Your care team may give you fluids and electrolytes with your treatment. Talk to your care team if you have: • Decrease in your amount of urine • Blood in your urine • Swelling of your ankles • Loss of appetite
Muscle or Joint Pain or Weakness	Description: Muscle pain is soreness, aching, cramps, stiffness, tenderness, or weakness in one or more muscles. Joint pain is pain, stiffness, swelling, or reduced movement where two bones meet. Recommendations: • Keep a pain diary: note pain levels, locations, and activities that make it better or worse. • Do gentle exercise (walking, stretching, yoga) to maintain mobility and strength. Check with your care team before starting a new activity. • Use a warm compress for stiff muscles or a cold pack to reduce swelling and numb pain—use what helps the area. • Your care team may recommend or prescribe medicines, including over-the-counter pain relievers. Talk to your care team if you have: • Pain you cannot control with usual measures • Swelling, redness, or warmth in a joint • New weakness • Trouble walking or moving

Hair Loss (Alopecia)	Description: Hair loss or thinning may begin days to weeks after treatment starts and usually grows back later. New growth can be a different color or texture and may not look the same as before. Recommendations: - Consider a short haircut before treatment and use scarves, hats, or wigs for comfort and confidence. - Keep your head covered outdoors to protect it from the sun and cold; use sunscreen on your uncovered scalp. - Use gentle haircare: mild shampoo, soft brush, and avoid heat styling and harsh treatments. - Ask your care team about wig prescriptions or resources for head coverings. Talk to your care team if you have: - No hair regrowth months after treatment ends - Concern about hair changes or need help finding a wig or support resources
----------------------	--

Select Rare Side Effects

Side Effect	Talk to Your Care Team if You Have Any of These Signs or Symptoms	
Changes in the Electrical Activity of Your Heart called QT prolongation	Feel faint, lightheaded, dizzy	Fast or irregular heartbeat
Lung Problems	- Cough - Shortness of breath	Chest pain
Severe Skin Reactions	- A red, itchy, skin rash - Peeling of your skin or blisters	- Severe itching - Fever of 100.4°F (38°C) or higher

**Before starting treatment, ask your care team when to call 9-1-1 or seek emergency help.
If you experience any new, worsening, or uncontrolled side effects, contact your care team immediately.**

Intimacy, Fertility, Pregnancy, and Breastfeeding

- Treatment may **change how you feel about intimacy and your body**. However, physical closeness—such as holding hands and hugging—remains safe. It is common to have questions about intimacy. If needed, talk to your care team for guidance.
- Treatment may affect your **ability to have children**. It may damage your reproductive organs or stop them from working. If you are worried about fertility, talk to your care team before starting treatment.
- Treatment may **harm an unborn baby**.
 - If you are able to become pregnant:
 - Take a pregnancy test before starting treatment.
 - Use an effective method of birth control during treatment and for 3 weeks after your last dose of ribociclib.
 - If you think you might be pregnant or if you become pregnant, tell your care team right away.
 - If your partner is able to become pregnant, use an effective method of birth control—such as condoms—during treatment with ribociclib.
- **Do NOT breastfeed** during treatment and for 3 weeks after your last dose of ribociclib.

Additional Information

- **Tell your care team about all the medicines you take.**
This includes prescriptions, over-the-counter drugs, vitamins, and herbal products. Before starting any new medicine, supplement, or vaccine, ask your care team first.
- **This Patient Education Sheet may not describe all possible side effects.**
Call your care team for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Notes

Updated Date: June 10, 2026

Scan the QR code below to access this education sheet.



Important notice: The Association of Cancer Care Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), Network for Collaborative Oncology Development & Advancement, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated in gathering information for and developing this patient education guide. This guide represents a brief summary of the medication derived from information provided by the drug manufacturer and other resources.

This guide does not cover all existing information related to the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this medication and should not substitute for the advice of a qualified healthcare professional. Provision of this guide is for informational purposes only and does not constitute or imply endorsement, recommendation, or favoring of this medication by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional.

Permission: Patient Education Sheets are provided as a free educational resource for patients with cancer and their caregivers in need of concise, easy to understand information about cancer therapy. Healthcare providers are permitted to copy and distribute the sheets to patients as well as direct patients to the Patient Education Sheets website. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holders. Permission requests, including direct linking from Electronic Health Records, and licensing inquiries should be emailed to patienteducationsheets@ncoda.org.

Copyright © 2026 by Network for Collaborative Oncology Development & Advancement, Inc. All rights reserved.

PES-197