

Inavolisib

Care Team Contact Information: _____

Pharmacy Contact Information: _____

Diagnosis: _____

- This treatment is used for breast cancer.
- It may also be used for other reasons.
- Your care team will perform a test for a specific type of abnormal endothelial growth factor receptor (EGFR) gene to make sure inavolisib is right for you.

Goal of Treatment: _____

- Treatment may continue for a certain time period, until it no longer works, or until side effects are no longer controlled.

Treatment Regimen

Treatment Name	How the Treatment Works	How the Treatment is Given
Inavolisib (IH-nuh-voh-LIH-sib); Itovebi (i-TOV-bee)	Slows down or stops the growth of cancer cells by blocking a specific protein that helps them survive.	Tablet(s) taken by mouth.

Treatment Administration and Schedule

Inavolisib is often given in combination with other treatments. Talk with your care team about your exact treatment and schedule.

Your inavolisib dosing instructions:

- Inavolisib comes in 2 tablet strengths: 3 mg and 9 mg.
- Your dose might differ, but inavolisib is typically taken as one 9 mg tablet by mouth once daily.
- Take inavolisib 1 time each day, at about the same time each day.
- Take inavolisib with or without food.
- Swallow inavolisib tablet(s) whole. Do not chew, crush, or split the tablet(s).
- Do not change your dose or stop taking inavolisib unless your care team tells you to.
- Your care team may tell you to change your dose, temporarily stop, or completely stop taking inavolisib if you develop certain side effects.
- If you miss a dose of inavolisib, you may still take your missed dose within 9 hours from the time you usually take it. If it has been more than 9 hours after you usually take your dose, skip the dose for that day. The next day, take the dose at your usual time.

Treatment Administration and Schedule (Continued)

- If you vomit after taking a dose of inavolisib, do not take an extra dose on that day. Take your regular dose of inavolisib at your usual time the next day.
- For women who have not reached menopause or are just starting menopause, your healthcare provider will prescribe a medicine called luteinizing hormone-releasing hormone (LHRH) agonist. For men, your healthcare provider may prescribe a LHRH agonist.
- If you take too much inavolisib, call your care team or go to the nearest hospital emergency room right away.

Storage and Handling of Inavolisib

- Store inavolisib at room temperature between 68°F and 77°F (20°C and 25°C).
- Store inavolisib in the original container and keep the bottle tightly closed to protect from moisture.
- Keep inavolisib and all medicines out of the reach of children and pets.
- Ask your care team how to safely throw away any unused inavolisib.

Appointments

Appointments may include regular check-ups with your care team, lab visits, and imaging tests. It's important to keep your appointments whenever you can. If you miss an appointment, call your care team as soon as possible to reschedule it.

Supportive Care to Prevent and Treat Side Effects

Description	Supportive Care Taken at Home
To help prevent or treat nausea and vomiting	
Other	

Common Side Effects

Side Effect	Important Information
Low White Blood Cell (WBC) Count (Neutropenia) and Increased Risk of Infection	Description: WBCs help protect your body from infections. A low WBC count increases your risk of infection. Recommendations: - Wash your hands often and bathe regularly. - Avoid crowded places and close contact with people who are sick. - Follow food safety and wound care advice from your care team. - Your care team may prescribe medicine to help your WBCs recover. Talk to your care team if you have: - Fever of 100.4°F (38°C) or higher - Chills - New or worsening cough or sore throat - Painful urination or signs of a urinary infection - Feeling much more tired than usual - Red, swollen, warm, or painful areas on the skin (possible skin infection)
Low Platelet Count (Thrombocytopenia)	Description: Platelets help your blood clot and wounds heal. A low platelet count increases your risk of bruising and bleeding. Recommendations: - Blow your nose gently and avoid picking it. - Brush your teeth gently with a soft toothbrush and keep good oral hygiene. - Use an electric razor for shaving and a nail file instead of nail clippers. - Avoid over-the-counter medicines that can increase bleeding risk (for example, nonsteroidal anti-inflammatory drugs (NSAIDs) like ibuprofen). - Talk with your care team or dentist before medical or dental procedures — you may need to pause treatment. Talk to your care team if you have: - A nosebleed lasting more than 5 minutes despite pressure - A cut that continues to bleed - Heavy gum bleeding when brushing or flossing - Sudden or severe headache - Blood in your urine or stool - Blood in your spit after coughing
Low Red Blood Cell (RBC) Count and Hemoglobin (Hgb) (Anemia)	Description: RBCs and Hgb carry oxygen to your body's tissues and remove carbon dioxide. Low RBC or Hgb (anemia) can make you feel weak or very tired, or may make you look pale. Recommendations: - Aim for 7 to 8 hours of sleep each night. - Do not drive, operate heavy machinery, or do other dangerous activities if you are very tired. - Balance activity and rest — stay as active as you can, but rest when needed. - Eat a balanced diet and follow any nutrition or supplement advice from your care team. Talk to your care team if you have: - Shortness of breath - Dizziness or fainting - Fast or irregular heartbeats - Sudden or severe headache

Fatigue	Description: Fatigue is a constant and sometimes strong feeling of tiredness. Recommendations: • Routine exercise can help reduce fatigue. Talk with your care team to find the right type and amount of activity for you. • Ask family and friends for help with daily tasks and for emotional support. • Try healthy ways to feel better, such as meditation, journaling, yoga, or guided imagery, to reduce anxiety and improve well-being. • Aim for 7 to 8 hours of sleep each night. Limit daytime naps to help you sleep better at night. • Do not drive, operate heavy machinery, or do other potentially dangerous activities if you are very tired. Talk to your care team if you have: • Tiredness that affects your daily life or prevents you from doing normal activities • Tiredness that does not get better with rest • Dizziness or weakness along with severe tiredness
Mouth Sores or Irritation (Mucositis or Stomatitis)	Description: Treatment can irritate the lining of the mouth. In some cases, this can cause redness, sores, pain, and swelling. Recommendations: • Rinse your mouth after meals and at bedtime; rinse more often if sores develop. • Brush your teeth gently with a soft toothbrush or use a cotton swab after meals. • Use a mild, non-alcohol mouth rinse at least 4 times daily (after meals and at bedtime). Example: 1/8 teaspoon salt + 1/4 teaspoon baking soda in 8 oz warm water. • Avoid acidic, hot, spicy, rough, or crunchy foods and drinks that can irritate your mouth. • Avoid tobacco, alcohol, and alcohol-based mouthwash. • Your care team may prescribe medicines or mouth treatments to help with pain and healing. Talk to your care team if you have: • Painful mouth sores or throat pain • Trouble eating or significant weight loss

Nausea and Vomiting	Description: Nausea is an uncomfortable feeling in your stomach or the need to throw up. You may or may not vomit. Recommendations: • Eat smaller, more frequent meals. • Avoid fatty, fried, spicy, or highly sweet foods. • Eat bland foods at room temperature and drink clear liquids. • If you vomit, start with small sips of water, broth, or other clear liquids. If these stay down, try soft foods (such as gelatin, plain cornstarch pudding, yogurt, strained soup, or strained cooked cereal) and gradually return to solid foods. • Your care team may prescribe medicine for these symptoms. Talk to your care team if you have: • Vomiting for more than 24 hours • Nonstop vomiting • Signs of dehydration (very thirsty, dry mouth, dizziness, or dark urine) • Blood in your vomit or vomit that looks like coffee grounds • Severe stomach pain that does not go away after vomiting
Liver Problems	Description: Treatment can cause liver injury. Your care team may check your liver with blood tests before and during treatment. Talk to your care team if you have: • Yellowing of your skin or the white part of your eyes (jaundice) • Severe nausea or vomiting • Pain on the right side of your stomach area (abdomen) • Dark, tea-colored urine • Bleeding or bruising more easily than normal
Diarrhea	Description: Diarrhea is loose, watery stools or more frequent bowel movements than usual. It can cause dehydration and weakness. Recommendations: • Keep track of how often you go to the bathroom each day. • Drink 8 to 10 glasses of water or other fluids daily, unless your care team tells you otherwise. • Eat small meals of mild, low-fiber foods (such as bananas, applesauce, potatoes, chicken, rice, and toast). • If you have diarrhea, avoid high-fiber foods (such as raw vegetables, fruits, and whole grains), gas-producing foods (such as broccoli and beans), dairy (such as milk and yogurt), and spicy, fried, or greasy foods. • Your care team may recommend an antidiarrheal medicine such as loperamide (Imodium). Talk to your care team if you have: • 4 or more bowel movements above your usual number in 24 hours • Dizziness or lightheadedness while having diarrhea • Signs of dehydration (very thirsty, dry mouth, dizziness, or dark urine) • Bloody diarrhea

Kidney Problems	Description: Treatment can cause kidney problems, including damage to the kidneys and decreased kidney function. Your care team will monitor your kidney function during treatment. Recommendations: • Drink 8 to 10 glasses of water or other fluids each day, unless your care team tells you otherwise. • Your care team may give you fluids and electrolytes with your treatment. Talk to your care team if you have: • Decrease in your amount of urine • Blood in your urine • Swelling of your ankles • Loss of appetite
High Blood Sugar (Hyperglycemia)	Description: Treatment may cause high blood sugar levels. Your care team may perform blood tests to check for these changes and treat you if needed. Recommendations: • Eat a well-balanced diet. • Limit sugary drinks and foods. • Eat smaller, more frequent meals. • Be physically active for at least 30 minutes most days. • Your care team may ask you to check your blood sugar at home. If you are already doing this, they may ask you to do it more frequently. Talk to your care team if you have: • Frequent urination • Drowsiness • Increased thirst • Loss of appetite • Blurred vision • Fruity smell on your breath • Confusion • Nausea, vomiting, or stomach pain • It becomes harder to control your blood sugar

Rash or Itchy Skin	Description: Rash or itchy skin can cause redness, swelling, and a variety of bumps or patches (small red spots, welts, blisters, or scaly dry areas). Recommendations: • Keep your skin moisturized with creams or lotions to reduce rash and itchiness. • Wear loose-fitting clothing. • Avoid perfumes and colognes, as they may worsen rash symptoms. • Limit time spent in the heat to prevent worsening symptoms. • Avoid sun exposure, especially between 10 AM and 4 PM, to lower the risk of sunburn. • Wear long-sleeved clothing with ultraviolet (UV) protection and broad-brimmed hats. • Apply broad-spectrum sunscreen (UVA/UVB) with sun protective factor (SPF) 30 or higher as directed. • Use lip balm with SPF 30 or higher. • Avoid tanning beds. • Your care team may recommend medicines for symptoms. Talk to your care team if you have: • Rash or itching that continues to worsen
Changes in Electrolytes and Other Laboratory Results	Description: Treatment may cause low levels of calcium, potassium, sodium, and magnesium in your blood. Your care team will perform blood tests to check for these changes and will treat you if needed. Talk to your care team if you have: • Tingling or numbness around the mouth, hands, or feet • New weakness or trouble breathing • Sudden weight gain • Swelling of your arms, hands, legs, or ankles • New or worsening muscle weakness, spasms, tremors, or cramps • Irregular, fast, or pounding heartbeat, or chest pain • Severe dizziness, fainting, or lightheadedness • Severe constipation or stomach-area (abdominal) pain • New or worsening confusion, drowsiness, or trouble thinking • Severe headache • Seizures • Sudden weakness, unsteadiness, or trouble walking • Difficulty breathing • Extreme fatigue • Numbness or tingling in your fingers or toes • Nausea, vomiting, or loss of appetite

Before starting treatment, ask your care team when to call 9-1-1 or seek emergency help.
If you experience any new, worsening, or uncontrolled side effects, contact your care team immediately.

Intimacy, Fertility, Pregnancy, and Breastfeeding

- Treatment may **change how you feel about intimacy and your body**. However, physical closeness—such as holding hands and hugging—remains safe. It is common to have questions about intimacy. If needed, talk to your care team for guidance.
- Treatment can affect your **ability to have children**. It may damage your reproductive organs or stop them from working. If you are worried about fertility, talk to your care team before starting treatment.
- Treatment may **harm an unborn baby**.
 - If you are able to become pregnant:
 - Your care team will verify your pregnancy status before treatment.
 - Use an effective method of birth control during treatment and for 1 week after your last dose of inavolisib.
 - If you think you might be pregnant or if you become pregnant, tell your care team right away.
 - If your partner is able to become pregnant, use an effective method of birth control—such as condoms—during treatment and for 1 week after your last dose of inavolisib.
- **Do not breastfeed** during treatment and for 1 week after your last dose of inavolisib.

Additional Information

- **Tell your care team about all the medicines you take.**
This includes prescriptions, over-the-counter drugs, vitamins, and herbal products. Before starting any new medicine, supplement, or vaccine, ask your care team first.
- **This Patient Education Sheet may not describe all possible side effects.**
Call your care team for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Notes

Updated Date: July 24, 2026

Scan the QR code below to access this education sheet.

**Important Notice:**

The Association of Cancer Care Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), Network for Collaborative Oncology Development & Advancement, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated to gather information and develop this patient education guide. This guide is a summary of the medication, based on information provided by the drug manufacturer and other sources.

This guide does not cover all information available on the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this medication. It should not be used as a substitute for the advice of a qualified healthcare professional. The provision of this guide is for informational purposes only. It does not constitute or imply endorsement, recommendation, or favoring of this medication by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. All decisions regarding this medication should be made with the guidance and direction of a qualified healthcare professional.

Permission:

Patient Education Sheets are provided as a free educational resource for patients with cancer and their caregivers in need of concise, easy-to-understand information about cancer therapy. Healthcare providers may copy and distribute the sheets to patients and direct them to the Patient Education Sheets website. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holders. Permission requests, including direct linking from Electronic Health Records, and licensing inquiries should be emailed to patienteducationsheets@ncoda.org.

Copyright © 2026 by Network for Collaborative Oncology Development & Advancement, Inc. All rights reserved.

PES-577