

# Tafasitamab, Rituximab, and Lenalidomide (Tafa-R<sup>2</sup>)

**Care Team Contact Information:** \_\_\_\_\_

**Pharmacy Contact Information:** \_\_\_\_\_

**Diagnosis:** \_\_\_\_\_

- This treatment is used for follicular lymphoma (FL).
- It may also be used for other reasons.

**Goal of Treatment:** \_\_\_\_\_

- Treatment may continue for a certain time period, until it no longer works, or until side effects are no longer controlled.

## Treatment Regimen

This treatment is often called “Tafa R-squared.” You may see this written as “Tafa-R<sup>2</sup>” or “Tafa-R2”:

- Tafa: **Tafasitamab**
- R<sup>2</sup>:
  - Lenalidomide (**Revlimid**)
  - **Rituximab**

| Treatment Name                                                                             | How the Treatment Works                                                                                 | How the Treatment is Given                        |
|--------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|---------------------------------------------------|
| Tafasitamab (TA-fuh-SIH-tuh-mab):<br>Monjuvi (mon-JOO-vee)                                 | Helps your immune system find and attack cancer cells by targeting a specific protein on their surface. | Infusion into a vein (intravenous (IV) infusion). |
| Rituximab (rih-TUK-sih-mab):<br>Rituxan (rih-TUK-sun), Riabni, Ruxience, Truxima, Zimrixby | Helps your immune system find and attack cancer cells by targeting a specific protein on their surface. | Infusion into a vein (intravenous (IV) infusion). |
| Lenalidomide (leh-nuh-LIH-doh-mide):<br>Revlimid (REV-lih-mid)                             | Boosts the immune system, cuts off the cancer's blood supply, and directly attacks the cancer cells.    | Capsule(s) taken by mouth.                        |

**Note:** Your care team may give you rituximab and hyaluronidase (Rituxan Hycela) instead of rituximab. Rituximab and hyaluronidase is given as an injection under the skin (subcutaneous injection) into the stomach area (abdomen) over 5 to 7 minutes.

## Treatment Administration and Schedule

Treatment is typically repeated every 28 days. This length of time is called a “cycle.” You may receive up to 5 cycles with Tafa-R<sup>2</sup>, followed by treatment with tafasitamab and lenalidomide for up to 7 cycles. Your care team will tell you how many treatment cycles you will receive.

## Treatment Administration and Schedule (Continued)

## Cycle 1

- Tafasitamab is given on Days 1, 8, 15, and 22.
- Rituximab is given Days 1, 8, 15, and 22.
- Lenalidomide is taken 1 time a day on Days 1 to 21, followed by 7 days off.

| Treatment Name                            | Cycle 1, Day(s) |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       |
|-------------------------------------------|-----------------|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|-----------------|-------|
|                                           | 1               | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22              | 23-28 |
| Treatment Given at the Clinic or Hospital |                 |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       |
| Tafasitamab                               | ✓               |   |   |   |   |   |   | ✓ |   |    |    |    |    |    | ✓  |    |    |    |    |    |    | ✓               |       |
| Rituximab                                 | ✓               |   |   |   |   |   |   | ✓ |   |    |    |    |    |    | ✓  |    |    |    |    |    |    | ✓               |       |
| Treatment Taken at Home                   |                 |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       |
| Lenalidomide                              | ✓               | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | Week-long break |       |

## Cycles 2 and 3

- Tafasitamab is given on Days 1, 8, 15, and 22.
- Rituximab is given on Day 1.
- Lenalidomide is taken 1 time a day on Days 1 to 21, followed by 7 days off.

| Treatment Name                            | Cycle 2, Day(s) |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       | Next Cycle |
|-------------------------------------------|-----------------|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|-----------------|-------|------------|
|                                           | 1               | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22              | 23-28 | Day 1      |
| Treatment Given at the Clinic or Hospital |                 |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       |            |
| Tafasitamab                               | ✓               |   |   |   |   |   |   | ✓ |   |    |    |    |    |    | ✓  |    |    |    |    |    |    | ✓               |       | ✓          |
| Rituximab                                 | ✓               |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       | ✓          |
| Treatment Taken at Home                   |                 |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       |            |
| Lenalidomide                              | ✓               | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | Week-long break |       | ✓          |

**Cycles 4 and 5**

- Tafasitamab is given on Days 1 and 15.
- Rituximab is given on Day 1.
- Lenalidomide is taken 1 time a day on Days 1 to 21, followed by 7 days off.

| Treatment Name                            | Cycle 4, Day(s) |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       | Next Cycle |
|-------------------------------------------|-----------------|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|-----------------|-------|------------|
|                                           | 1               | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22              | 23-28 | Day 1      |
| Treatment Given at the Clinic or Hospital |                 |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       |            |
| Tafasitamab                               | ✓               |   |   |   |   |   |   |   |   |    |    |    |    |    | ✓  |    |    |    |    |    |    |                 |       | ✓          |
| Rituximab                                 | ✓               |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       | ✓          |
| Treatment Taken at Home                   |                 |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       |            |
| Lenalidomide                              | ✓               | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | Week-long break |       | ✓          |

**Cycles 6 to 12**

- Tafasitamab is given on Days 1 and 15.
- Lenalidomide is taken 1 time a day on Days 1 to 21, followed by 7 days off.

| Treatment Name                            | Cycle 6, Day(s) |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       | Next Cycle |
|-------------------------------------------|-----------------|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|-----------------|-------|------------|
|                                           | 1               | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22              | 23-28 | Day 1      |
| Treatment Given at the Clinic or Hospital |                 |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       |            |
| Tafasitamab                               | ✓               |   |   |   |   |   |   |   |   |    |    |    |    |    | ✓  |    |    |    |    |    |    |                 |       | ✓          |
| Treatment Taken at Home                   |                 |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |                 |       |            |
| Lenalidomide                              | ✓               | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | ✓  | Week-long break |       | ✓          |

**Lenalidomide Information**

Your lenalidomide dosing instructions:

- Lenalidomide comes in 6 capsule strengths: 2.5 mg, 5 mg, 10 mg, 15 mg, 20 mg, and 25 mg.
- Your dose might differ, but lenalidomide is typically taken as one 20 mg capsule by mouth once daily for 21 days (3 weeks), followed by 7 days (1 week) off.
- Take lenalidomide 1 time each day, around the same time each day.
- Swallow lenalidomide capsules whole with water. Do not open, break, or chew your capsules.
- Lenalidomide may be taken with or without food.
- Do not change your dose or stop taking lenalidomide unless your care team tells you to.
- Your care team may tell you to change your dose, temporarily stop, or completely stop taking lenalidomide if you develop certain side effects.
- If you vomit or miss a dose of lenalidomide and it has been less than 12 hours since your regular time, take it as soon as you remember. If it has been more than 12 hours, just skip your missed dose. Do not take 2 doses at the same time.
- If you take too much lenalidomide, call your care team or go to the nearest hospital emergency room right away.

**Storage and Handling of Lenalidomide**

- Store lenalidomide at room temperature between 68°F and 77°F in a dry location away from light.
- Keep lenalidomide out of the reach of children and pets.
- Whenever possible, give lenalidomide to yourself and follow the steps below. If someone else gives it to you, they must also follow these steps:
  1. Wash hands with soap and water.
  2. Put on gloves to avoid touching the medication. Note: Gloves are not needed if you give the drug to yourself.
  3. Transfer the lenalidomide from its package to a small medicine or other disposable cup.
  4. Administer the medicine immediately by mouth with water.
  5. Remove gloves, if used, and throw them and medicine cup in household trash.
  6. Wash hands with soap and water.
- Do not open or break lenalidomide capsules or handle them any more than needed.
  - If powder from the lenalidomide capsule comes in contact with your skin, wash the skin right away with soap and water.
  - If powder from the lenalidomide capsule comes in contact with the inside of your eyes, nose, or mouth, flush well with water.
- If you plan to use a daily pill box or pill reminder, contact your care team before using it.
  - When the box or reminder is empty, wash it with soap and water before refilling.
  - The person refilling the box or reminder should:
- Wear gloves. Note: Gloves are not needed if you are refilling it yourself.
- Wash their hands with soap and water after completing the task, regardless of whether gloves were worn.
- Ask your care team how to safely throw away any unused lenalidomide. Do not throw it in the trash or flush it down the sink or toilet.

### Appointments

Appointments may include regular check-ups with your care team, treatment appointments, lab visits, and imaging tests. It's important to keep your appointments whenever you can. If you miss an appointment, call your care team as soon as possible to reschedule it.

### Supportive Care to Prevent and Treat Side Effects

| Description                                          | Supportive Care Given at the Clinic or Hospital | Supportive Care Taken at Home |
|------------------------------------------------------|-------------------------------------------------|-------------------------------|
| To help lower the risk of infections                 |                                                 |                               |
| To help lower the risk of blood clots                |                                                 |                               |
| To help prevent or treat nausea and vomiting         |                                                 |                               |
| To help lower the risk of infusion-related reactions |                                                 |                               |
| Other                                                |                                                 |                               |

## Common Side Effects

| Side Effect                                                                                           | Important Information                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Low White Blood Cell (WBC) Count (Neutropenia) and Increased Risk of Infection (Boxed Warning)</b> | <p>Description: WBCs help protect your body from infections. A low WBC count increases your risk of infection.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>Wash your hands often and bathe regularly.</li> <li>Avoid crowded places and close contact with people who are sick.</li> <li>Follow food safety and wound care advice from your care team.</li> <li>Your care team may prescribe medicine to help your WBCs recover.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>Fever of 100.4°F (38°C) or higher</li> <li>Chills</li> <li>New or worsening cough or sore throat</li> <li>Painful urination or signs of a urinary infection</li> <li>Feeling much more tired than usual</li> <li>Red, swollen, warm, or painful areas on the skin (possible skin infection)</li> </ul>                                                                                                                                                                                                                             |
| <b>Low Platelet Count (Thrombocytopenia) (Boxed Warning)</b>                                          | <p>Description: Platelets help your blood clot and wounds heal. A low platelet count increases your risk of bruising and bleeding.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>Blow your nose gently and avoid picking it.</li> <li>Brush your teeth gently with a soft toothbrush and keep good oral hygiene.</li> <li>Use an electric razor for shaving and a nail file instead of nail clippers.</li> <li>Avoid over-the-counter medicines that can increase bleeding risk (for example, nonsteroidal anti-inflammatory drugs (NSAIDs) like ibuprofen).</li> <li>Talk with your care team or dentist before medical or dental procedures — you may need to pause treatment.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>A nosebleed lasting more than 5 minutes despite pressure</li> <li>A cut that continues to bleed</li> <li>Heavy gum bleeding when brushing or flossing</li> <li>Sudden or severe headache</li> <li>Blood in your urine or stool</li> <li>Blood in your spit after coughing</li> </ul> |
| <b>Low Red Blood Cell (RBC) Count and Hemoglobin (Hgb) (Anemia)</b>                                   | <p>Description: RBCs and Hgb carry oxygen to your body's tissues and remove carbon dioxide. Low RBC or Hgb (anemia) can make you feel weak or very tired, or may make you look pale.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>Aim for 7 to 8 hours of sleep each night.</li> <li>Do not drive, operate heavy machinery, or do other dangerous activities if you are very tired.</li> <li>Balance activity and rest — stay as active as you can, but rest when needed.</li> <li>Eat a balanced diet and follow any nutrition or supplement advice from your care team.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>Shortness of breath</li> <li>Dizziness or fainting</li> <li>Fast or irregular heartbeats</li> <li>Sudden or severe headache</li> </ul>                                                                                                                                                                                                                                                     |

|                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|----------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Fatigue        | <p>Description: Fatigue is a constant and sometimes strong feeling of tiredness.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>• Routine exercise can help reduce fatigue. Talk with your care team to find the right type and amount of activity for you.</li> <li>• Ask family and friends for help with daily tasks and for emotional support.</li> <li>• Try healthy ways to feel better, such as meditation, journaling, yoga, or guided imagery, to reduce anxiety and improve well-being.</li> <li>• Aim for 7 to 8 hours of sleep each night. Limit daytime naps to help you sleep better at night.</li> <li>• Do not drive, operate heavy machinery, or do other potentially dangerous activities if you are very tired.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>• Tiredness that affects your daily life or prevents you from doing normal activities</li> <li>• Tiredness that does not get better with rest</li> <li>• Dizziness or weakness along with severe tiredness</li> </ul>                                                                                                                    |
| Liver Problems | <p>Description: Treatment can cause liver injury. Your care team may check your liver with blood tests before and during treatment.</p> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>• Yellowing of your skin or the white part of your eyes (jaundice)</li> <li>• Severe nausea or vomiting</li> <li>• Pain on the right side of your stomach area (abdomen)</li> <li>• Dark, tea-colored urine</li> <li>• Bleeding or bruising more easily than normal</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Diarrhea       | <p>Description: Diarrhea is loose, watery stools or more frequent bowel movements than usual. It can cause dehydration and weakness.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>• Keep track of how often you go to the bathroom each day.</li> <li>• Drink 8 to 10 glasses of water or other fluids daily, unless your care team tells you otherwise.</li> <li>• Eat small meals of mild, low-fiber foods (such as bananas, applesauce, potatoes, chicken, rice, and toast).</li> <li>• If you have diarrhea, avoid high-fiber foods (such as raw vegetables, fruits, and whole grains), gas-producing foods (such as broccoli and beans), dairy (such as milk and yogurt), and spicy, fried, or greasy foods.</li> <li>• Your care team may recommend an antidiarrheal medicine such as loperamide (Imodium).</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>• 4 or more bowel movements above your usual number in 24 hours</li> <li>• Dizziness or lightheadedness while having diarrhea</li> <li>• Signs of dehydration (very thirsty, dry mouth, dizziness, or dark urine)</li> <li>• Bloody diarrhea</li> </ul> |

|                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|--------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Constipation       | <p>Description: Constipation means hard, dry stools or fewer bowel movements than normal. It can cause discomfort or pain.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>• Track how often you have bowel movements each day.</li> <li>• Drink 8 to 10 glasses of fluids daily, unless your care team says otherwise.</li> <li>• Stay active and exercise regularly.</li> <li>• Eat more high-fiber foods (such as raw fruits, vegetables, and whole grains) unless advised otherwise.</li> <li>• Your care team may recommend laxatives such as polyethylene glycol (MiraLAX) or senna.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>• Constipation lasting 3 or more days</li> <li>• No bowel movement 48 hours after using a laxative</li> </ul>                                                                                                                                                                                                                                                                                               |
| Rash or Itchy Skin | <p>Description: Rash or itchy skin can cause redness, swelling, and a variety of bumps or patches (small red spots, welts, blisters, or scaly dry areas).</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>• Keep your skin moisturized with creams or lotions to reduce rash and itchiness.</li> <li>• Wear loose-fitting clothing.</li> <li>• Avoid perfumes and colognes, as they may worsen rash symptoms.</li> <li>• Limit time spent in the heat to prevent worsening symptoms.</li> <li>• Avoid sun exposure, especially between 10 AM and 4 PM, to lower the risk of sunburn.</li> <li>• Wear long-sleeved clothing with ultraviolet (UV) protection and broad-brimmed hats.</li> <li>• Apply broad-spectrum sunscreen (UVA/UVB) with sun protective factor (SPF) 30 or higher as directed.</li> <li>• Use lip balm with SPF 30 or higher.</li> <li>• Avoid tanning beds.</li> <li>• Your care team may recommend medicines for symptoms.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>• Rash or itching that continues to worsen</li> </ul> |

## Select Rare Side Effects

| Side Effect                                                             | Talk to Your Care Team if You Have Any of These Signs or Symptoms                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |                                                                                                                                                                                                                                                                                                     |                                                                                                                                                                                                                                                                          |
|-------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Blood Clots (Boxed Warning)</b>                                      | <p>Signs or symptoms of a <b>blood clot</b> in the lung, arm, or leg may include:</p> <ul style="list-style-type: none"> <li>• Shortness of breath</li> <li>• Chest pain</li> <li>• Arm or leg swelling</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                   | <p>Signs and symptoms of a <b>heart attack</b> may include:</p> <ul style="list-style-type: none"> <li>• Chest pain that may spread to the arms, neck, jaw, back, or stomach area (abdomen)</li> <li>• Feeling sweaty</li> <li>• Shortness of breath</li> <li>• Feeling sick or vomiting</li> </ul> | <p>Signs or symptoms of a <b>stroke</b> may include:</p> <ul style="list-style-type: none"> <li>• Sudden numbness or weakness, especially on one side of the body</li> <li>• Severe headache or confusion</li> <li>• Problems with vision, speech, or balance</li> </ul> |
| <b>Severe Skin and Mouth Reactions (Boxed Warning)</b>                  | <ul style="list-style-type: none"> <li>• Painful sores or ulcers on your skin, lips or in your mouth</li> <li>• Blisters</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | <ul style="list-style-type: none"> <li>• Peeling skin</li> <li>• Rash</li> <li>• Pustules</li> </ul>                                                                                                                                                                                                |                                                                                                                                                                                                                                                                          |
| <b>Hepatitis B Virus (HBV) Reactivation (Boxed Warning)</b>             | <p>Before you receive rituximab, your care team will do blood tests to check for HBV infection. If you have had hepatitis B or are a carrier of hepatitis B virus, receiving rituximab could cause the virus to become an active infection again. Hepatitis B reactivation may cause serious liver problems, including liver failure and death. Your care team will monitor you for hepatitis B infection during and for several months after you stop receiving rituximab.</p> <ul style="list-style-type: none"> <li>• Worsening tiredness</li> <li>• Yellowing of your skin or white part of your eyes</li> </ul> |                                                                                                                                                                                                                                                                                                     |                                                                                                                                                                                                                                                                          |
| <b>Progressive Multifocal Leukoencephalopathy (PML) (Boxed Warning)</b> | <p>PML is a rare, serious brain infection caused by a virus that can happen in people who receive rituximab. People with weakened immune systems can get PML. PML can result in death or severe disability. There is no known treatment, prevention, or cure for PML.</p> <ul style="list-style-type: none"> <li>• Confusion</li> <li>• Dizziness or loss of balance</li> <li>• Difficulty walking or talking</li> <li>• Decreased strength or weakness on one side of your body</li> <li>• Vision problems, such as blurred vision or loss of vision</li> </ul>                                                     |                                                                                                                                                                                                                                                                                                     |                                                                                                                                                                                                                                                                          |
| <b>Infusion-Related Reactions (Boxed Warning)</b>                       | <p>Your care team may slow or temporarily stop the infusion if you have a reaction. Treatment may be permanently stopped for a severe reaction.</p> <ul style="list-style-type: none"> <li>• Chills or shaking</li> <li>• Itching, rash, or flushing</li> <li>• Trouble breathing, wheezing, or tongue swelling</li> <li>• Dizziness or feeling faint</li> <li>• Feeling of impending doom</li> <li>• Fever of 100.4°F (38°C) or higher</li> <li>• New or severe pain in your back or neck</li> </ul>                                                                                                                |                                                                                                                                                                                                                                                                                                     |                                                                                                                                                                                                                                                                          |

|                                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |                                                                                                                                                                                                                                                                               |
|------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Heart Problems                     | <ul style="list-style-type: none"> <li>• Fast or irregular heartbeats</li> <li>• Swelling of your stomach area (abdomen), legs, hands, feet, or ankles</li> <li>• Shortness of breath</li> <li>• New or worsening chest discomfort, including pain or pressure</li> </ul>                                                                                                                                                                                                                                                                 | <ul style="list-style-type: none"> <li>• Nausea or vomiting</li> <li>• Weight gain</li> <li>• Pain or discomfort in your arms, back, neck, or jaw</li> <li>• Protruding neck veins</li> <li>• Breaking out in a cold sweat</li> <li>• Feeling lightheaded or dizzy</li> </ul> |
| Serious Infections                 | <p>Fatal and serious infections, including opportunistic infections, have occurred in patients treated with tafasitamab and rituximab.</p> <ul style="list-style-type: none"> <li>• Fever of 100.4°F (38°C) or higher</li> <li>• Severe cough with or without an increase in mucus (sputum) production</li> <li>• Severe sore throat</li> <li>• Shortness of breath</li> <li>• Burning or pain when urinating</li> <li>• Unusual vaginal discharge or irritation</li> <li>• Redness, swelling, or pain in any part of the body</li> </ul> |                                                                                                                                                                                                                                                                               |
| Stomach and Serious Bowel Problems | <ul style="list-style-type: none"> <li>• Severe stomach-area (abdomen) pain</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                    | <ul style="list-style-type: none"> <li>• Repeated vomiting during treatment</li> </ul>                                                                                                                                                                                        |
| Kidney Problems                    | <ul style="list-style-type: none"> <li>• Decrease in your amount of urine</li> <li>• Blood in your urine</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                       | <ul style="list-style-type: none"> <li>• Swelling of your ankles</li> <li>• Loss of appetite</li> </ul>                                                                                                                                                                       |
| Tumor Lysis Syndrome (TLS)         | <p>Tumor lysis happens when cancer cells break apart and release large amounts of substances into your bloodstream faster than your body can handle. TLS is a group of conditions that affect your heart, kidneys, and muscles.</p> <ul style="list-style-type: none"> <li>• Severe nausea, vomiting, or diarrhea</li> <li>• Urinating smaller amounts or having dark-colored urine</li> <li>• Muscle cramps or twitching</li> <li>• Rapid heartbeats or chest pain</li> <li>• Confusion or weakness</li> <li>• Seizures</li> </ul>       |                                                                                                                                                                                                                                                                               |
| Thyroid Gland Problems             | <ul style="list-style-type: none"> <li>• Extreme tiredness</li> <li>• Feeling hot or cold</li> <li>• Your voice gets deeper</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                    | <ul style="list-style-type: none"> <li>• Weight gain or loss</li> <li>• Hair loss</li> </ul>                                                                                                                                                                                  |
| Injection-Site Reactions           | <ul style="list-style-type: none"> <li>• Itching</li> <li>• Swelling</li> <li>• Bruising or bleeding</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                           | <ul style="list-style-type: none"> <li>• Pain</li> <li>• Rash or redness of the skin</li> </ul>                                                                                                                                                                               |
| Risk of New Cancers                | <p>There is a risk of developing one or more new cancers during or after treatment. Talk with your care team about this risk, and ask about the signs and symptoms to watch for.</p>                                                                                                                                                                                                                                                                                                                                                      |                                                                                                                                                                                                                                                                               |

**Before starting treatment, ask your care team when to call 9-1-1 or seek emergency help.**  
**If you experience any new, worsening, or uncontrolled side effects, contact your care team immediately.**

### Intimacy, Pregnancy, and Breastfeeding

- Treatment may **change how you feel about intimacy and your body**. However, physical closeness—such as holding hands and hugging—remains safe. It is common to have questions about intimacy. If needed, talk to your care team for guidance.
- **Boxed Warning:** Treatment may cause **birth defects or death of an unborn baby**. If you are pregnant or plan to become pregnant, you must not take lenalidomide.
  - You must not get pregnant:
    - For at least 4 weeks before starting lenalidomide
    - While taking lenalidomide
    - During any breaks (interruptions) in your treatment with lenalidomide
    - For at least 4 weeks after stopping lenalidomide
  - If you can become pregnant:
    - You will have pregnancy tests weekly for 4 weeks, then every 4 weeks if your menstrual cycle is regular, or every 2 weeks if your menstrual cycle is irregular.
    - If you miss your period or have unusual bleeding, you will need to have a pregnancy test and receive counseling.
    - Must agree to use two acceptable forms of birth control at the same time, for at least 4 weeks before, while taking, during any breaks (interruptions) in your treatment, and for at least 4 weeks after stopping lenalidomide.
    - Talk with your care team to find out about options for acceptable forms of birth control that you may use to prevent pregnancy before, during, and after treatment with lenalidomide.
    - Use an effective method of birth control during treatment with Tafa-R<sup>2</sup>, for 3 months after your last dose of tafasitamab, and for 12 months after your last dose of rituximab.
  - If your partner is able to become pregnant, use an effective method of birth control—such as condoms—during treatment with Tafa-R<sup>2</sup>
  - If you become pregnant while taking lenalidomide, stop taking it right away and call your care team. If your care team is not available, you can call the REMS Call Center at 1-888-423-5436. Healthcare providers and patients should report all cases of pregnancy to:
    - FDA MedWatch at 1-800-FDA-1088, and
    - REMS Call Center at 1-888-423-5436
- A pregnancy exposure registry collects information about pregnancies that occur when a patient takes lenalidomide or when a pregnant partner may have been exposed through semen. Call the Lenalidomide REMS program for more information.

**Intimacy, Pregnancy, and Breastfeeding (Continued)**

- Lenalidomide can pass into human semen:
  - Even if you have had a vasectomy, you must always use a latex or synthetic condom during any sexual contact with anyone who is or can become pregnant while taking lenalidomide, during any breaks (interruptions) in your treatment with lenalidomide, and for up to 4 weeks after stopping lenalidomide.
  - Do not have unprotected sexual contact with anyone who is or could become pregnant. Tell your care team if you do have unprotected sexual contact with anyone who is or could become pregnant.
  - Do not donate sperm while taking lenalidomide, during any breaks (interruptions) in your treatment, and for up to 4 weeks after stopping lenalidomide. If anyone becomes pregnant with your sperm, the baby may be exposed to lenalidomide and may be born with birth defects.
- If your partner becomes pregnant, you should call your healthcare provider right away.
- **Do not breastfeed** during treatment with Tafa-R<sup>2</sup>, for 3 months after your last dose of tafasitamab, and for 6 months after your last dose of rituximab.

**Handling Body Fluids and Waste**

Some drugs you receive may stay in your urine, stool, sweat, or vomit for many days after treatment. Because many cancer drugs are toxic, your body waste may also be dangerous to touch. To help protect yourself, your loved ones, and the environment, **follow these instructions** for at least **48 hours** after each dose of **lenalidomide**: (Note: Tafasitamab and rituximab do not require special handling of body fluids and waste.)

- People who are pregnant should avoid touching anything that may be soiled with body fluids from the patient.
- You can use your usual toilet. Always close the lid and flush to discard all waste. If you have a low-flow toilet, flush twice.
- If the toilet or seat is soiled with urine, stool, or vomit, clean the surface after each use before others use it.
- Wash your hands with soap and water for at least 20 seconds after using the toilet.
- If you need a bedpan, inform your caregiver so they can wear gloves and assist with cleanup. Wash the bedpan with soap and water daily.
- If you cannot control your bladder or bowels, use a disposable pad with a plastic back, a diaper, or a sheet to absorb waste.
- Wash any skin exposed to body waste with soap and water.
- Wash soiled linens or clothing separately from other laundry. If you don't have a washer, place them in a plastic bag until they can be washed.
- Wash your hands with soap and water after touching soiled linens or clothing.

**Additional Information**

- **Tell your care team about all the medicines you take.**  
This includes prescriptions, over-the-counter drugs, vitamins, and herbal products. Before starting any new medicine, supplement, or vaccine, ask your care team first.
- **You should not receive a “live vaccine”** before, during, or after treatment, until your care team tells you it is okay. If you are not sure about the type of immunization or vaccine, ask your care team. These vaccines may not be safe or may not work as well during treatment.
- **Do not donate blood** while you take lenalidomide, during any breaks (interruptions) in your treatment, and for 4 weeks after stopping lenalidomide. If someone who is pregnant gets your donated blood, their baby may be exposed to lenalidomide and may be born with birth defects.
- **Before starting lenalidomide, read and agree to the Lenalidomide REMS program** and sign the Patient-Physician Agreement Form. For details, call 1-888-423-5436 or visit [www.lenalidomiderems.com](http://www.lenalidomiderems.com).
- Lenalidomide cannot be automatically refilled. You must complete the required REMS steps for each prescription.
- **This Patient Education Sheet may not describe all possible side effects.**  
Call your care team for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

**Notes**

Updated Date: September 14, 2026

Scan the QR code below to access this education sheet.

**Important Notice:**

The Association of Cancer Care Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), Network for Collaborative Oncology Development & Advancement, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated to gather information and develop this patient education guide. This guide is a summary of the medication, based on information provided by the drug manufacturer and other sources.

This guide does not cover all information available on the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this medication. It should not be used as a substitute for the advice of a qualified healthcare professional. The provision of this guide is for informational purposes only. It does not constitute or imply endorsement, recommendation, or favoring of this medication by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. All decisions regarding this medication should be made with the guidance and direction of a qualified healthcare professional.

**Permission:**

Patient Education Sheets are provided as a free educational resource for patients with cancer and their caregivers in need of concise, easy-to-understand information about cancer therapy. Healthcare providers may copy and distribute the sheets to patients and direct them to the Patient Education Sheets website. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holders. Permission requests, including direct linking from Electronic Health Records, and licensing inquiries should be emailed to [patienteducationsheets@ncoda.org](mailto:patienteducationsheets@ncoda.org).

Copyright © 2026 by Network for Collaborative Oncology Development & Advancement, Inc. All rights reserved.

PES-682