

Pertuzumab, Paclitaxel, Carboplatin, and Trastuzumab (P-TCH)

Care Team Contact Information: _____

Pharmacy Contact Information: _____

Diagnosis: _____

- This treatment is used for breast cancer.
- It may also be used for other reasons.
- Your care team will perform a test for a specific type of abnormal human epidermal growth factor receptor 2 (HER2) gene to make sure treatment is right for you.

Goal of Treatment: _____

- Treatment may continue for a certain time period, until it no longer works, or until side effects are no longer controlled.

Treatment Regimen

This treatment is often called P-TCH, PTCH, or TCHP:

- **P:** Pertuzumab
- **T:** Taxol (Paclitaxel)
- **C:** Carboplatin
- **H:** Herceptin (Trastuzumab)

Treatment Name	How the Treatment Works	How the Treatment is Given
Pertuzumab (per-TOO-zoo-mab): Perjeta (per-JEH-tuh), Poherdy	Slows down or stops cancer growth by blocking a specific protein that cancer cells need to survive.	Infusion into a vein (intravenous (IV) infusion).
Paclitaxel (PA-klih-TAK-sil): Taxol (TAK-ol)	Slows down or stops the growth of cancer cells by preventing cancer cells from properly dividing and creating new cells.	Infusion into a vein (intravenous (IV) infusion).
Carboplatin (KAR-boh-pla-tin): Paraplatin (PAIR-ah-PLAT-in)	Slows down or stops the growth of cancer cells by damaging the genetic material that cancer cells need to multiply.	Infusion into a vein (intravenous (IV) infusion).
Trastuzumab (tras-TOO-zoo-mab): Herceptin (her-SEP-tin), Hercessi, Herzuma, Kanjinti, Ogivri, Ontruzant, Trazimera	Slows down or stops cancer growth by blocking a specific protein that cancer cells need to survive.	Infusion into a vein (intravenous (IV) infusion).

Note: Your care team may give you trastuzumab and hyaluronidase (Herceptin Hylecta) instead of trastuzumab. Trastuzumab and hyaluronidase is given as an injection under the skin (subcutaneous injection) into the thigh over 2 to 5 minutes.

Treatment Administration and Schedule

Treatment is typically repeated every 21 days. This length of time is called a “cycle.” You may receive up to 9 cycles with P-TCH followed by treatment with pertuzumab and trastuzumab alone. Your care team will tell you how many treatment cycles you will receive and how long you will continue taking trastuzumab.

- Pertuzumab is given on Day 1.
- Paclitaxel is given on Day 1 and 8.
- Carboplatin is given on Day 1.
- Trastuzumab is given on Day 1.

Treatment Name											Next Cycle
	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Day 8	...	Day 21	Day 1
Pertuzumab	✓										✓
Paclitaxel	✓							✓			✓
Carboplatin	✓										✓
Trastuzumab	✓										✓

Appointments

Appointments may include regular check-ups with your care team, treatment appointments, lab visits, and imaging tests. It's important to keep your appointments whenever you can. If you miss an appointment, call your care team as soon as possible to reschedule it.

Supportive Care to Prevent and Treat Side Effects

Description	Supportive Care Given at the Clinic or Hospital	Supportive Care Taken at Home
To help your body make white blood cells to fight infections		
To help prevent or treat nausea and vomiting		
To help lower the risk of infusion-related reactions		
To help lower the risk of numbness, tingling, or pain in the hands or feet (peripheral neuropathy)		
Other		

Common Side Effects

Side Effect	Important Information
Low White Blood Cell (WBC) Count (Neutropenia) and Increased Risk of Infection (Boxed Warning)	Description: WBCs help protect your body from infections. A low WBC count increases your risk of infection. Recommendations: - Wash your hands often and bathe regularly. - Avoid crowded places and close contact with people who are sick. - Follow food safety and wound care advice from your care team. - Your care team may prescribe medicine to help your WBCs recover. Talk to your care team if you have: - Fever of 100.4°F (38°C) or higher - Chills - New or worsening cough or sore throat - Painful urination or signs of a urinary infection - Feeling much more tired than usual - Red, swollen, warm, or painful areas on the skin
Low Platelet Count (Thrombocytopenia) (Boxed Warning)	Description: Platelets help your blood clot and wounds heal. A low platelet count increases your risk of bruising and bleeding. Recommendations: - Blow your nose gently and avoid picking it. - Brush your teeth gently with a soft toothbrush and keep good oral hygiene. - Use an electric razor for shaving and a nail file instead of nail clippers. - Avoid over-the-counter medicines that can increase bleeding risk (for example, nonsteroidal anti-inflammatory drugs (NSAIDs) like ibuprofen). - Talk with your care team or dentist before medical or dental procedures — you may need to pause treatment. Talk to your care team if you have: - A nosebleed lasting more than 5 minutes despite pressure - A cut that continues to bleed - Heavy gum bleeding when brushing or flossing - Sudden or severe headache - Blood in your urine or stool - Blood in your spit after coughing
Low Red Blood Cell (RBC) Count and Hemoglobin (Hgb) (Anemia) (Boxed Warning)	Description: Red blood cells carry oxygen throughout your body. Low levels can make you feel tired, weak, dizzy, or short of breath. Recommendations: - Aim for 7 to 8 hours of sleep each night. - Do not drive, operate heavy machinery, or do other dangerous activities if you are very tired. - Balance activity and rest — stay as active as you can, but rest when needed. - Eat a balanced diet and follow any nutrition or supplement advice from your care team. Talk to your care team if you have: - Shortness of breath - Dizziness or fainting - Fast or irregular heartbeats - Sudden or severe headache

Nausea and Vomiting (Boxed Warning)	Description: Nausea is an uncomfortable feeling in your stomach or the need to throw up. You may or may not vomit. Recommendations: • Eat smaller, more frequent meals. • Avoid fatty, fried, spicy, or highly sweet foods. • Eat bland foods at room temperature and drink clear liquids. • If you vomit, start with small sips of water, broth, or other clear liquids. If these stay down, try soft foods (such as gelatin, plain cornstarch pudding, yogurt, strained soup, or strained cooked cereal) and gradually return to solid foods. • Your care team may prescribe medicine for these symptoms. Talk to your care team if you have: • Vomiting for more than 24 hours • Nonstop vomiting • Signs of dehydration (very thirsty, dry mouth, dizziness, or dark urine) • Blood in your vomit or vomit that looks like coffee grounds • Severe stomach pain that does not go away after vomiting
Allergic Reactions, Including Anaphylaxis (Boxed Warning)	Description: Allergic reactions are common during treatment with paclitaxel and carboplatin, and can sometimes be serious. Serious allergic reactions can happen during or after your paclitaxel and carboplatin infusion, including life-threatening allergic reactions. Talk to your care team if you have: • Itchy, raised bumps on the skin (hives) • Coughing that does not go away • Nausea or vomiting • Stomach (abdominal) pain • Increased saliva • Breathing problems such as wheezing • Throat tightness or change in voice • Fever of 100.4°F or higher • Chest discomfort • Chills or shaking • Back pain

Fatigue	Description: Fatigue is a constant and sometimes strong feeling of tiredness. Recommendations: • Routine exercise can help reduce fatigue. Talk with your care team to find the right type and amount of activity for you. • Ask family and friends for help with daily tasks and for emotional support. • Try healthy ways to feel better, such as meditation, journaling, yoga, or guided imagery, to reduce anxiety and improve well-being. • Aim for 7 to 8 hours of sleep each night. Limit daytime naps to help you sleep better at night. • Do not drive, operate heavy machinery, or do other potentially dangerous activities if you are very tired. Talk to your care team if you have: • Tiredness that affects your daily life or prevents you from doing normal activities • Tiredness that does not get better with rest • Dizziness or weakness along with severe tiredness
Diarrhea	Description: Diarrhea means having bowel movements more often than usual. Your stools may also be loose or watery. If you have diarrhea: • Keep track of how often you have diarrhea. • Drink 8 to 10 glasses of water or fluids each day. Do this unless your care team tells you otherwise. • Eat small meals with low-fiber foods. Try bananas, applesauce, potatoes, chicken, rice, or toast. • Avoid high-fiber foods. These include raw vegetables, fruit, and whole grains. • Avoid foods that may cause gas. These include broccoli and beans. • Avoid milk, yogurt, and other dairy foods. • Avoid spicy, fried, or greasy foods. • Your care team may recommend medicine for diarrhea. One example is loperamide (Imodium). Talk to your care team if you have: • An increase of 4 or more bowel movements than usual in one day • Dizziness or lightheadedness while you have diarrhea • Signs of dehydration, such as feeling very thirsty, dizzy, or having dark urine • Blood in your stool

Headache	Description: Headaches are pain in your head that can vary in intensity, frequency, and location. Recommendations: • Keep a headache diary to track the frequency, duration, intensity, and triggers of your headaches. • Stay hydrated by drinking plenty of water, as dehydration can contribute to headaches. • Apply a cold or warm compress to your forehead or neck to help ease headache pain. • Get adequate sleep (7 to 8 hours per night) and establish a regular sleep schedule. • Limit caffeine intake. • Your care team may recommend medicine for headaches. Talk to your care team if you have: • Severe headache • More frequent or worsening headaches • Dizziness or feeling faint • Confusion • Changes in vision
Numbness, Tingling, or Burning in Your Hands or Feet (Peripheral Neuropathy)	Description: Nerve pain and tingling are uncomfortable sensations from nerve damage or irritation. Pain may be sharp, burning, or deep. Tingling can feel like pins-and-needles or mild electric shocks, often in the hands, feet, arms, or legs. Recommendations: • Keep a daily log of pain and sensations, noting triggers and what helps or makes it worse. • Check your feet every day for cuts, sores, blisters, or color changes, especially if numbness reduces feeling. • Wear comfortable, well-fitting shoes and avoid walking barefoot if sensation is reduced. • Protect hands and feet from extreme heat or cold. • Your care team may recommend or prescribe medicines, topical treatments, physical therapy, or supplements to help with symptoms. Talk to your care team if you have: • New or worsening “pins and needles”, burning, or numbness in your hands or feet • Trouble moving your arms or legs, or weakness • Problems with balance or frequent falls

Hair Loss (Alopecia)	Description: Hair loss or thinning may begin days to weeks after treatment starts and usually grows back later. New growth can be a different color or texture and may not look the same as before. Recommendations: - Consider a short haircut before treatment and use scarves, hats, or wigs for comfort and confidence. - Keep your head covered outdoors to protect it from the sun and cold; apply sunscreen to your exposed scalp. - Use gentle haircare: mild shampoo, soft brush, and avoid heat styling and harsh treatments. - Ask your care team about wig prescriptions or resources for head coverings. Talk to your care team if you have: - No hair regrowth months after treatment ends - Concern about hair changes or need help finding a wig or support resources
----------------------	--

Select Rare Side Effects

Side Effect	Talk to Your Care Team if You Have Any of These Signs or Symptoms
Heart Problems (Boxed Warning)	- Fast or irregular heartbeats - Swelling of your stomach area (abdomen), legs, hands, feet, or ankles - Shortness of breath - Nausea or vomiting - New or worsening chest discomfort, including pain or pressure - Weight gain - Pain or discomfort in your arms, back, neck, or jaw - Protruding neck veins - Breaking out in a cold sweat - Feeling lightheaded or dizzy
Lung Problems (Boxed Warning)	Trastuzumab can cause lung problems that may be serious, life-threatening, or fatal. Lung problems can include inflammation or scarring in the lungs called interstitial lung disease (ILD) or pneumonitis. If you develop lung problems, your care team may treat you with corticosteroid medicines. Call your care team right away if you develop: - Cough - Trouble breathing or shortness of breath - Fever - Other new or worsening breathing symptoms such as chest tightness or wheezing
Infusion-Related Reactions (Boxed Warning)	Your care team may slow or temporarily stop the infusion if you have a reaction. Treatment may be permanently stopped for a severe reaction. - Chills or shaking - Itching, rash, or flushing - Trouble breathing, wheezing, or tongue swelling - Dizziness or feeling faint - Feeling of impending doom - Fever of 100.4°F (38°C) or higher - New or severe pain in your back or neck

Liver Problems	• Yellowing of your skin or the white part of your eyes (jaundice) • Severe nausea or vomiting	• Pain on the right side of your stomach area (abdomen) • Dark, tea-colored urine • Bleeding or bruising more easily than normal
Kidney Problems	• Decrease in your amount of urine • Blood in your urine	• Swelling of your ankles • Loss of appetite
Eye Problems	• Dry or red eyes • Eye pain or swelling • Vision changes	• Increased tears • Sensitivity to light • Blurred vision
Hearing Loss	• New or worsening hearing loss • Ringing, buzzing, or other noises in your ears (tinnitus)	• Trouble understanding speech or needing higher volume on devices • Dizziness or balance problems

Before starting treatment, ask your care team when to call 9-1-1 or seek emergency help.
If you experience any new, worsening, or uncontrolled side effects, contact your care team immediately.

Intimacy, Fertility, Pregnancy, and Breastfeeding

- Treatment may **change how you feel about intimacy and your body**. However, physical closeness—such as holding hands and hugging—remains safe. It is common to have questions about intimacy. If needed, talk to your care team for guidance.
- Treatment can affect your **ability to have children**. It may damage your reproductive organs or stop them from working. If you are worried about fertility, talk to your care team before starting treatment.
- **Boxed Warning:** Treatment may **harm an unborn baby**.
 - If you are able to become pregnant:
 - Your care team will verify your pregnancy status before treatment.
 - Use an effective method of birth control during treatment with P-TCH, for 3 to 6 months after your last dose of carboplatin, and for 7 months after your last dose of trastuzumab and pertuzumab.
 - If you think you might be pregnant or if you become pregnant, tell your care team right away.
 - If your partner is able to become pregnant, use an effective method of birth control—such as condoms—during treatment with P-TCH and for 3 to 6 months after your last dose of carboplatin.
- **Do not breastfeed** during treatment with P-TCH, for 6 to 10 days after your last dose of paclitaxel, and for 7 months after your last dose of trastuzumab and pertuzumab.

Handling Body Fluids and Waste

Some drugs you receive may stay in your urine, stool, sweat, or vomit for many days after treatment. Because many cancer drugs are toxic, your body waste may also be dangerous to touch. To help protect yourself, your loved ones, and the environment, **follow these instructions** for at least **48 hours** after each dose of **carboplatin** and for at least **4 days** after each dose of **paclitaxel**: (Note: Pertuzumab and trastuzumab do not require special handling of body fluids and waste.)

- People who are pregnant should avoid touching anything that may be soiled with body fluids from the patient.
- You can use your usual toilet. Always close the lid and flush to discard all waste. If you have a low-flow toilet, flush twice.
- If the toilet or seat is soiled with urine, stool, or vomit, clean the surface after each use before others use it.
- Wash your hands with soap and water for at least 20 seconds after using the toilet.
- If you need a bedpan, inform your caregiver so they can wear gloves and assist with cleanup. Wash the bedpan with soap and water daily.
- If you cannot control your bladder or bowels, use a disposable pad with a plastic back, a diaper, or a sheet to absorb waste.
- Wash any skin exposed to body waste with soap and water.
- Wash soiled linens or clothing separately from other laundry. If you don't have a washer, place them in a plastic bag until they can be washed.
- Wash your hands with soap and water after touching soiled linens or clothing.

Additional Information

- **Tell your care team about all the medicines you take.**
This includes prescriptions, over-the-counter drugs, vitamins, and herbal products. Before starting any new medicine, supplement, or vaccine, ask your care team first.
- **This Patient Education Sheet may not describe all possible side effects.**
Call your care team for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Notes

Updated Date: October 12, 2026

Scan the QR code below to access this education sheet.

**Important Notice:**

The Association of Cancer Care Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), Network for Collaborative Oncology Development & Advancement, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated to gather information and develop this patient education guide. This guide is a summary of the medication, based on information provided by the drug manufacturer and other sources.

This guide does not cover all information available on the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this medication. It should not be used as a substitute for the advice of a qualified healthcare professional. The provision of this guide is for informational purposes only. It does not constitute or imply endorsement, recommendation, or favoring of this medication by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. All decisions regarding this medication should be made with the guidance and direction of a qualified healthcare professional.

Permission:

Patient Education Sheets are provided as a free educational resource for patients with cancer and their caregivers in need of concise, easy-to-understand information about cancer therapy. Healthcare providers may copy and distribute the sheets to patients and direct them to the Patient Education Sheets website. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holders. Permission requests, including direct linking from Electronic Health Records, and licensing inquiries should be emailed to patienteducationsheets@ncoda.org.

Copyright © 2026 by Network for Collaborative Oncology Development & Advancement, Inc. All rights reserved.

PES-695